

Seminariefrågor - Cancergenetik och tumörbiologi

1. Vad menas med en mutation? Vad skiljer en konstitutionell mutation från en somatisk mutation?
2. Hur kan man klassificera cancerna utifrån deras funktioner?
3. Identifiera två muterade stabilitetsgener som förklarar en stor andel av ärftligheten vid bröstcancer och kolorektalcancer. Vilken funktion har stabilitetsgener ("caretakers")?
4. Vad kännetecknar ärftliga cancersyndrom? Vad menar man med att patienter med konstitutionella mutationer i cancerner har ett försprång ("head start") i tumörutvecklingen?
5. Definiera en gatekeeper-gen och ge några väl valda exempel på sådana gener. Spelar ordningen då mutationer uppstår någon roll?
6. Har tumörceller en högre mutationsfrekvens än normala celler?
7. Vad innebär kromosomal instabilitet (CIN)?
8. Vad menar författarna egentligen med "pathway"? Finns det någon möjlighet att experimentellt avgöra om två gener är i "samma pathway"?
9. Vilka är de tre huvudtyperna av signalvägar som är muterade i cancer och på vilka sätt bidrar de till tumörbildning?
10. Hur kan man använda kunskapen om signalvägar för diagnostik och läkemedelsutveckling?
11. Vad kan heterogenitet i tumören tänkas ha för konsekvenser för diagnostik och behandling?
12. I de flesta solida tumörer är inaktiverande mutationer i tumör-suppressorgener vanligare än aktiverande mutationer i onkgener. Varför är det ett problem ur läkemedelssynpunkt?
13. Olika tumörformer har väldigt olika förekomst av mutationer i genomet. Varför hittar man färre mutationer i t.ex. barntumörer?
14. Finns det mutationer som uppträder specifikt i metastaser?

Cancer genes and the pathways they control

Bert Vogelstein & Kenneth W Kinzler

The revolution in cancer research can be summed up in a single sentence: cancer is, in essence, a genetic disease. In the last decade, many important genes responsible for the genesis of various cancers have been discovered, their mutations precisely identified, and the pathways through which they act characterized. The purposes of this review are to highlight examples of progress in these areas, indicate where knowledge is scarce and point out fertile grounds for future investigation.

What we know

The cast. Alterations in three types of genes are responsible for tumorigenesis: oncogenes, tumor-suppressor genes and stability genes (Tables 1 and 2). Unlike diseases such as cystic fibrosis or muscular dystrophy, wherein mutations in one gene can cause disease, no single gene defect 'causes' cancer. Mammalian cells have multiple safeguards to protect them against the potentially lethal effects of cancer gene mutations, and only when several genes are defective does an invasive cancer develop. Thus it is best to think of mutated cancer genes as contributing to, rather than causing, cancer.

Oncogenes are mutated in ways that render the gene constitutively active or active under conditions in which the wild-type gene is not. Oncogene activations can result from chromosomal translocations, from gene amplifications or from subtle intragenic mutations affecting crucial residues that regulate the activity of the gene product. For example, the most common activating mutation of *BRAF* in human cancers changes a valine to a glutamate at codon 599, a residue within the activation loop of the kinase domain¹. The activation loop is normally regulated by phosphorylation at adjacent residues (Thr598 and Ser601). This suggests that the glutamate substitution at codon 599 mimics a phosphate group and constitutively activates the enzyme even in the absence of signals that would normally result in phosphorylation of the adjacent threonine or serine residues. The activated *BRAF* kinase then phosphorylates downstream targets such as extracellular signal-regulated kinase (ERK), leading to aberrant growth²(Fig. 1). A mutation in an oncogene is analogous to a stuck accelerator in an automobile; the car still moves forward even when the driver removes his foot from it. An activating somatic mutation in one allele of an oncogene is generally sufficient to confer a selective growth advantage on the cell.

Tumor-suppressor genes are targeted in the opposite way by genetic alterations: mutations reduce the activity of the gene

product. Such inactivations arise from missense mutations at residues that are essential for its activity, from mutations that result in a truncated protein, from deletions or insertions of various sizes, or from epigenetic silencing. A mutation in a tumor-suppressor gene is analogous to a dysfunctional brake in an automobile; the car doesn't stop even when the driver attempts to engage it. Some recently described tumor-suppressor genes have been hypothesized to exert a selective advantage on a cell when only one allele is inactivated and the other remains functional (that is, haploinsufficiency)³. However, mutations in both the maternal and paternal alleles of a tumor-suppressor gene are generally required to confer a selective advantage to the cell. This situation commonly arises through the deletion of one allele via a gross chromosomal event—such as loss of an entire chromosome or chromosome arm—coupled with an intragenic mutation of the other allele⁴.

Oncogene and tumor-suppressor gene mutations all operate similarly at the physiologic level: they drive the neoplastic process by increasing tumor cell number through the stimulation of cell birth or the inhibition of cell death or cell-cycle arrest. The increase can be caused by activating genes that drive the cell cycle, by inhibiting normal apoptotic processes or by facilitating the provision of nutrients through enhanced angiogenesis (Figs. 2–4). A third class of cancer genes, called stability genes or caretakers, promotes tumorigenesis in a completely different way when mutated. This class includes the mismatch repair (MMR), nucleotide-excision repair (NER) and base-excision repair (BER) genes responsible for repairing subtle mistakes made during normal DNA replication or induced by exposure to mutagens (Table 1). Other stability genes control processes involving large portions of chromosomes, such as those responsible for mitotic recombination and chromosomal segregation (for example, *BRCA1*, *BLM* and *ATM*; Table 1). Stability genes keep genetic alterations to a minimum, and thus when they are inactivated, mutations in other genes occur at a higher rate⁵. All genes are potentially affected by the resultant increased rate of mutation, but only mutations in oncogenes and tumor-suppressor genes affect net cell growth and can thereby confer a selective growth advantage to the mutant cell. As with tumor-suppressor genes, both alleles of stability genes generally must be inactivated for a physiologic effect to result.

The authors are at the Howard Hughes Medical Institute and The Sidney Kimmel Comprehensive Cancer Center, The Johns Hopkins University Medical Institutions, Baltimore, Maryland 21231, USA.
e-mail: vogelbe@welch.jhu.edu or kinzlike@jhmi.edu

Published online 30 July 2004; doi:10.1038/nm1087




Figure 1 Receptor tyrosine kinase (RTK) pathway. Gene products in red boxes indicate that the corresponding genes are mutated in the germline (and generally also are somatically mutated in nonfamilial tumors; see **Table 1**). Gene products in green boxes indicate that the corresponding gene has been found to be mutated only somatically (see **Table 2**). ‘GPG’ denotes growth-promoting-genes—that is, genes that stimulate cell proliferation or inhibit the rate of cell death or arrest. Diamonds (◆) indicate protein-protein interactions. Red arrows and red T-bars indicate transcriptional induction and repression, respectively. Small circled ‘P’, ‘OH’ and ‘Ub’ represent covalently attached phosphate, hydroxyl and ubiquitin groups, respectively.

In the analogy to autos, stability genes represent the mechanics and a defective stability gene is akin to an inept mechanic.

Mutations in these three classes of genes can occur in the germline, resulting in hereditary predispositions to cancer (**Table 1**), or in single somatic cells, resulting in sporadic tumors (**Tables 1 and 2**). It is important to point out that a mutation is defined as any change in the sequence of the genome. These changes include those affecting single base pairs as well as those creating large or small deletions or insertions, amplifications or translocations. In the germline, the most common mutations are subtle (point mutations or small deletions or insertions), whereas all types of mutation can be found in tumor cells. In fact, cancers represent one of the few disease types in which somatic mutations occurring after birth are pathogenic.

The first somatic mutation in an oncogene or tumor-suppressor gene that causes a clonal expansion initiates the neoplastic process⁶. Subsequent somatic mutations result in additional rounds of clonal expansion (and thus in tumor progression)⁷. Indeed, the best modern definition of a neoplastic cell is one that has clonally expanded as a result of somatic mutations^{6,7}. Germline mutations of these genes cause cancer predisposition, not cancer *per se*: people with these mutations have a ‘head start’ on the neoplastic process, as a mutation that can contribute to cancer is already present in every one of their cells. Such individuals therefore often develop multiple tumors that occur at an earlier age than in individuals whose cancer-gene mutations have all occurred

somatically⁴. Examples of hereditary syndromes associated with inherited mutations are listed in **Table 1**. In people with these syndromes, only a very small fraction of the total cells in an at-risk organ become neoplastic because other (somatic) mutations are required to develop a clinically detectable lesion. In nearly all dominantly inherited syndromes caused by tumor-suppressor genes and stability genes, the first somatic mutation affects the normal copy of the gene inherited from the unaffected parent⁴. Interestingly, the most common forms of hereditary cancer predisposition, leading to breast and colon cancers, are caused by inherited mutations of stability genes rather than tumor-suppressor genes or oncogenes (**Table 1**).

The plot. As noted above, cancer-gene mutations enhance net cell growth. As a result of research performed over the past decade, it is clear that there are many fewer pathways than genes. This concept is very familiar to geneticists studying yeast, flies, mice or worms—there are almost always a variety of genes that, when altered, lead to similar phenotypes^{8,9}. Application of this concept to cancer has been solidified by elucidation of the biochemical functions of the altered cancer genes, either in cell culture systems, in mice or in other organisms. For example, several cancer genes directly control transitions from a resting stage (G0 or G1) to a replicating phase (S) of the cell cycle (**Fig. 2a**, Rb pathway). The products of these genes include proteins as diverse as cdk4 (a kinase), cyclin D1 (which interacts with and activates cdk4), Rb (a transcription factor) and p16 (which interacts with and inhibits cdk4)^{10–12}. The genes encoding Rb and p16 are tumor-suppressor genes inactivated by mutation, whereas those encoding cdk4 and cyclin D1 are oncogenes activated by mutation. In addition to functional studies in model systems, detailed studies of individual tumor types have also provided compelling evidence that these four genes function in a single pathway in human cancers. Such studies have shown that the mutations within this pathway obey an ‘exclusivity principle’: that is, one and only one of the four genes noted above is generally mutated in any single tumor, exactly as predicted if the functional effect of each mutation was similar^{10–13}.

Another example of the reason for focusing on pathways rather than individual genes has been provided by studies of the *TP53* tumor-suppressor gene. The p53 protein is a transcription factor that normally inhibits cell growth and stimulates cell death when induced by cellular stress^{14–16}. The most common way to disrupt the p53 pathway is through a point mutation that inactivates its capacity to bind specifically to its cognate recognition sequence. However, there are several other ways to achieve the same effect, including amplification of the *MDM2* gene and infection with DNA tumor viruses whose products (such as the E6 protein of human papilloma virus) bind to p53 and functionally inactivate it (**Fig. 2b**, p53 pathway).

One of the most important—and most curious—discoveries of the 1990s was that virtually all DNA tumor viruses that cause tumors in experimental animals or humans encode proteins that inactivate both Rb and p53 (refs. 17–19). Of the hundreds of cancer genes known or remaining to be discovered, why should these two have been singled out as targets for inactivation by all DNA tumor viruses? The answer may be that it is impossible for a tumor of epithelial origin to form unless the p53 and Rb tumor-suppressor gene pathways have been inactivated. This conjecture is supported by studies showing that these two pathways are altered in a large fraction of many types of cancers. We predict that most of the cancers that now appear to be devoid of mutations in these two

© 2004 Nature Publishing Group <http://www.nature.com/naturemedicine>



Table 1 Cancer predisposition genes

Gene (synonym(s)) ^a	Syndrome	Hereditary pattern	Second hit	Pathway ^b	Major heredity tumor types ^c
Tumor-suppressor genes					
<i>APC</i>	FAP	Dominant	Inactivation of WT allele	APC	Colon, thyroid, stomach, intestine
<i>AXIN2</i>	Attenuated polyposis	Dominant	Inactivation of WT allele	APC	Colon
<i>CDH1</i> (E-cadherin)	Familial gastric carcinoma	Dominant	Inactivation of WT allele	APC	Stomach
<i>GPC3</i>	Simpson-Golabi-Behmel syndrome	X-linked	?	APC	Embryonal
<i>CYLD</i>	Familial cylindromatosis	Dominant	Inactivation of WT allele	APOP	Pilomatricomas
<i>EXT1,2</i>	Hereditary multiple exostoses	Dominant	Inactivation of WT allele	GLI	Bone
<i>PTCH</i>	Gorlin syndrome	Dominant	Inactivation of WT allele	GLI	Skin, medulloblastoma
<i>SUFU</i>	Medulloblastoma predisposition	Dominant	Inactivation of WT allele	GLI	Skin, medulloblastoma
<i>FH</i>	Hereditary leiomyomatosis	Dominant	Inactivation of WT allele	HIF1	Leiomyomas
<i>SDHB, C, D</i>	Familial paraganglioma	Dominant	Inactivation of WT allele	HIF1	Paragangliomas, pheochromocytomas
<i>VHL</i>	Von Hippel–Lindau syndrome	Dominant	Inactivation of WT allele	HIF1	Kidney
<i>TP53</i> (<i>p53</i>)	Li-Fraumeni syndrome	Dominant	Inactivation of WT allele	p53	Breast, sarcoma, adrenal, brain...
<i>WT1</i>	Familial Wilms tumor	Dominant	Inactivation of WT allele	p53	Wilms ^d
<i>STK11</i> (<i>LKB1</i>)	Peutz-Jeghers syndrome	Dominant	Inactivation of WT allele	PI3K	Intestinal, ovarian, pancreatic
<i>PTEN</i>	Cowden syndrome	Dominant	Inactivation of WT allele	PI3K	Hamartoma, glioma, uterus
<i>TSC1, TSC2</i>	Tuberous sclerosis	Dominant	Inactivation of WT allele	PI3K	Hamartoma, kidney
<i>CDKN2A</i> (<i>p16^{INK4A}, p14^{ARF}</i>)	Familial malignant melanoma	Dominant	Inactivation of WT allele	RB	Melanoma, pancreas
<i>CDK4</i>	Familial malignant melanoma	Dominant	?	RB	Melanoma
<i>RB1</i>	Hereditary retinoblastoma	Dominant	Inactivation of WT allele	RB	Eye
<i>NF1</i>	Neurofibromatosis type 1	Dominant	Inactivation of WT allele	RTK	Neurofibroma
<i>BMPR1A</i>	Juvenile polyposis	Dominant	Inactivation of WT allele	SMAD	Gastrointestinal
<i>MEN1</i>	Multiple endocrine neoplasia type I	Dominant	Inactivation of WT allele	SMAD	Parathyroid, pituitary, islet cell, carcinoid
<i>SMAD4</i> (<i>DPC4</i>)	Juvenile polyposis	Dominant	Inactivation of WT allele	SMAD	Gastrointestinal
<i>BHD</i>	Birt-Hogg-Dube syndrome	Dominant	Inactivation of WT allele	?	Renal, hair follicle
<i>HRPT2</i>	Hyperparathyroidism Jaw-tumor syndrome.	Dominant	Inactivation of WT allele	?	Parathyroid, jaw fibroma
<i>NF2</i>	Neurofibromatosis type 2	Dominant	Inactivation of WT allele	?	Meningioma, acoustic neuroma
Stability genes					
<i>MUTYH</i>	Attenuated polyposis	Recessive	?	BER	Colon
<i>ATM</i>	Ataxia telangiectasia	Recessive	?	CIN	Leukemias, lymphomas, brain
<i>BLM</i>	Bloom syndrome	Recessive	?	CIN	Leukemias, lymphomas, skin
<i>BRCA1, BRCA2</i>	Hereditary breast cancer	Dominant	Inactivation of WT allele	CIN	Breast, ovary
<i>FANCA, C, D2, E, F, G</i>	Fanconi anemia	Recessive	?	CIN	Leukemias
<i>NBS1</i>	Nijmegen breakage syndrome	Recessive	?	CIN	Lymphomas, brain
<i>RECQL4</i>	Rothmund-Thomson syndrome	Recessive	?	CIN	Bone, skin
<i>WRN</i>	Werner syndrome	Recessive	?	CIN	Bone, brain
<i>MSH2, MLH1, MSH6, PMS2</i>	HNPCC	Dominant	Inactivation of WT allele	MMR	Colon, uterus
<i>XPA, C; ERCC2–5; DDB2</i>	Xeroderma pigmentosum	Recessive	?	NER	Skin
Oncogenes					
<i>KIT</i>	Familial gastrointestinal stromal tumors	Dominant	?	RTK	Gastrointestinal stromal tumors
<i>MET</i>	Hereditary papillary renal cell carcinoma	Dominant	Mutant allele duplication	RTK	Kidney
<i>PDGFRA</i>	Familial gastrointestinal stromal tumors	Dominant	?	RTK	Gastrointestinal stromal tumors
<i>RET</i>	Multiple endocrine neoplasia type II	Dominant	Mutant allele duplication	RTK	Thyroid, parathyroid, adrenal

WT, wild type. ^aRepresentative genes of all the major pathways and hereditary cancer predisposition types are listed. For a complete list, see ref. 117. Approved gene symbols are provided for each entry, with alternative names in parentheses. ^bIn many cases, the gene has been implicated in several pathways. The single pathway that is listed for each gene represents a 'best guess' (when one can be made) and for the reasons noted in the text and in the legend to **Figure 9**, should not be regarded as conclusive. APOP, apoptotic pathway; RTK, receptor tyrosine kinase pathway (see **Fig. 1**). ^cIn most cases, the nonfamilial tumor spectrum caused by somatic mutations of the gene includes those occurring in the familial cases plus additional tumor types. For example, mutations of *TP53* and *CDKN2A* are found in many more tumor types than those to which Li-Fraumeni and familial malignant melanoma patients, respectively, are predisposed.

© 2004 Nature Publishing Group <http://www.nature.com/naturemedicine> npg




Figure 2 Rb (a) and p53 (b) pathways. Symbols as in Figure 1.

pathways will eventually be shown to contain them. This prediction will become testable once more efficient methods of detecting mutations are developed and all the genes in the pathways become known.

In addition to the Rb and p53 pathways, there are others that have a role in many tumor types, including those involving adenomatous polyposis coil (APC), glioma-associated oncogene (GLI), hypoxia-inducible transcription factor (HIF)-1, phosphoinositide 3-kinase (PI3K), SMADs and receptor tyrosine kinases (RTKs) (Figs. 1 and 4–8). In each case, mutations in multiple members of the pathway have been found to be mutated in more than one type of cancer, and in many cases the mutations within a single pathway obey the exclusivity principle noted above. The discoveries of these signal transduction pathways represent major scientific success stories of the last decade²⁰. Their delineation has practical as well as theoretical implications. For example, they predict that targeted therapies directed against a particular gene product may be active against a tumor with a mutation of the targeted gene or any gene upstream of the target. Additionally, the fact that defects in a relatively small number of pathways underlie many different tumor types suggests that targeted therapeutics will be effective against a broad range of cancers.

Tumors can be broadly classified as liquid or solid. The former includes leukemias and lymphomas, composed of neoplastic cells whose precursors are normally mobile. Solid tumors are composed of epithelial or mesenchymal cells that normally are immobile. There are numerous other differences between liquid and solid tumors. For example, at least three mutations seem to be required to develop a malignant solid tumor in adults²¹; each of these mutations likely alters one of the pathways described in Figure 9. In contrast, only one or two mutations may be required to develop a malignant liquid tumor⁶. Perhaps liquid tumors don't require as many pathways to be

inactivated because their precursor cells are already mobile and invasive, key characteristics that solid tumor cells must develop to become malignant. Additionally, oncogene activations caused by chromosome translocation events are the most common genetic alterations observed in liquid tumors²². Consistent chromosome translocations are much less common in solid tumors, whereas inactivations of tumor-suppressor genes are ubiquitous²³. Finally, there are several genes that are uniquely altered in specific subtypes of liquid tumors, seem primarily to affect differentiation, do not obviously participate in the pathways depicted in Figure 9 and do not occur in hereditary form (Table 2). These molecular distinctions add to the cytogenetic, epidemiologic and medical evidence that liquid and solid tumors should be considered separately in

terms of their biology, behavior and pathogenesis. As solid tumors are much more common than liquid tumors, most of the examples provided in this review focus on the former.

Supporting cast. That solid tumors are composed of two compartments, one consisting of neoplastic epithelial cells and the other of stromal cells, was pointed out a hundred years ago²⁴. The importance of the interactions between stroma and epithelium is becoming increasingly recognized^{25,26}. And four discoveries made during the last ten years have propelled one component of the stroma, endothelial cells, into the spotlight.

First, naturally occurring inhibitors of angiogenesis were identified and shown to hinder the growth of experimental tumors^{27,28}. Second, synthetic inhibitors of one of the major regulatory pathways of angiogenesis (VEGF) were produced and shown to inhibit




Figure 3 Apoptosis pathway. Symbols as in Figure 1.

tumor progression in patients with cancer²⁹. Third, inactivation of *VHL*, a classic tumor-suppressor gene, was shown to support renal cell tumor growth through the control of tumor angiogenesis³⁰. The protein encoded by *VHL* is part of a ubiquitin ligase complex that degrades HIF-1 α in the presence of oxygen (Fig. 4, HIF1 pathway). In the absence of oxygen under normal circumstances or when *VHL* is mutated in tumors, the HIF-1 α transcription factor is stabilized, leading to the expression of cytokines such as VEGF and culminating in angiogenesis³¹. And finally, all oncogenes and tumor-suppressor gene pathways have been implicated in angiogenesis, either directly or indirectly, as illustrated by the central position of the HIF1 pathway in Figure 9. An understanding of tumor angiogenesis, blood flow, oxygenation and related issues involving the tumor-host relationship is becoming essential to studies of cancer biology as well as to the design of more effective forms of cancer therapy.

What we don't know

The opening act. The process of tumorigenesis is initiated when a replication-competent cell (stem cell or partially differentiated descendent of a stem cell) acquires a mutation in a 'gatekeeping' pathway that endows it with a selective growth advantage. In some cancers, the gatekeeper has been identified (examples are *RBI*, *APC* and *NFI* in tumors of the eye, colon and nervous system, respectively). In most common tumors, however, the gatekeeper is not known. It is also not known whether cancers of the lung, breast, prostate, bladder or brain can each be initiated through any one of several gatekeeping pathways or through only one. Many of the known gatekeepers were identified through the study of unusual families with predispositions to specific types of cancers⁴. There may be other families that provide clues to the nature of the gatekeepers in the future. We predict, however, that many of these remaining gatekeeping genes will only be identified through more 'brute-force' approaches involving sequence determination of major portions of the cancer cell genome³². As gatekeeping mutations provide fundamental insights into the biology and pathogenesis of particular cancers and are of singular importance to future diagnostic and therapeutic strategies, further research on this topic should be a priority.

The producers. As noted above, it appears that the cells of solid tumors must accumulate several rate-limiting mutations in cancer




Figure 4 HIF1 pathway. Symbols as in Figure 1.

genes to achieve malignant status. If these mutations had to occur simultaneously in a single cell, then the prevalence of cancer would be minimal³³. The current doctrine is that these mutations occur over time, with each mutation engendering a clonal expansion resulting in a large number of cells that then form a substrate for subsequent mutations⁶. Are normal rates of mutation, coupled with clonal expansions, sufficient to account for the prevalence of cancer, or is some form of genetic instability required for a cell to undergo these multiple, sequential mutations? This issue has been hotly debated for a long time^{34,35}, but the last 10 years of research have yielded some facts that clarify the issues. First, genetic instability can clearly contribute to cancer, as cogently demonstrated by the hereditary cancer predispositions caused by defects in stability genes (Table 1). Second, most cancers do not have a high mutation rate when this rate is measured at the nucleotide level. Accordingly, the number of mutations in a typical cancer cell is ~1 per megabase of DNA, similar to what would be expected in a normal cell that had passed through as many generations and population bottlenecks³⁶. These observations argue against a common role for defects in MMR, NER or BER in nonhereditary tumor types. However, there is another form of instability—chromosomal instability (CIN)—that is much more commonly found in cancers, and this is observed at the gross chromosomal rather than the nucleotide level³⁷. Though the actual rate of chromosomal changes has only been studied in a small number of cases, the end result of CIN, aneuploidy, is observed in nearly all solid tumors³⁸. At the molecular level, chromosome losses are evident as losses of heterozygosity (LOH). An average of 25–30% of the alleles present in normal cells are lost in cancers, and it is not unusual to observe losses of over 75% of the cell's alleles³⁷. Classic as well as modern cytogenetic studies are fully consistent with these observations^{23,39}. Such wholesale changes in chromosomal content can be advantageous for the cancer cell, allowing the efficient elimination of one allele of a tumor-suppressor gene as well as the production of variants that can rapidly adapt to changing microenvironments. The mechanisms underlying aneuploidy and chromosomal instability in sporadic tumors are still largely unknown, though some candidate genes and pathways, such as those involving cell-cycle checkpoints, telomere crisis or centrosomes, have been proposed^{34,40–44}. The role of telomeres in instability is particularly intriguing given the potential importance of telomerase in aging and the age-dependent incidence of most cancers^{42,45}. As genetic instabilities




Figure 5 APC pathway. Symbols as in Figure 1.

© 2004 Nature Publishing Group http://www.nature.com/naturemedicine






Figure 6 GLI pathway. Symbols as in Figure 1.

not only seem to be central to the neoplastic development but also may underlie the development of resistance to chemotherapeutic agents^{33,34}, the identification of the molecular mechanisms responsible for them is an important area of study.

The ending. Although abnormalities of the cancer genes listed in Tables 1 and 2 are essential contributors to cancer, most abnormalities in these genes occur relatively early in the disease process and none are known to be specifically associated with the metastatic stage. It is this final stage—the seeding and growth of satellite lesions in other organs—that is ultimately responsible for the great majority of neoplastic deaths⁴⁶. Primary tumors can generally be removed through surgery but widely metastatic lesions cannot be excised and are difficult or impossible to treat with adjuvant therapies.

Some of the biochemical processes involved in the early stages of metastasis, such as increased cell motility and production of matrix-degrading proteases, have been well studied⁴⁷. However, the genetic alterations responsible for endowing cells with these abilities have not been clearly identified. In fact, it has been suggested that metastasis is not dependent at all on new genetic abnormalities that occur after tumors have been established, and that the propensity to metastasize is determined early in the neoplastic process rather than near its end (reviewed in ref. 48). This suggestion is not readily compatible with the evidence that cancer is a genetic disease in which evolution occurs somatically. Just as macroevolution never stops, evolution of the cancer cell does not stop and new variants of tumor cells with potentially greater capacities to invade and metastasize are always being born. This evolution is itself driven by inherent genetic instabilities, as described above. However, this genetic perspective on metastasis cannot be validated—or invalidated—until a better understanding of the metastatic process is in hand.

Same actors in different roles. One might have expected that a specific mutation of a widely expressed gene would have identical or at least similar effects in different mammalian cell types. But this is not in general what is observed. Different effects of the same mutation are not only found in distinct cell types; differences can even be observed in the same cell type, depending on when the

mutation occurred during the tumorigenic process. *RAS* gene mutations provide informative examples of these complexities.

i. Cell type specificity: *KRAS2* gene mutations in normal pancreatic duct cells seem to initiate the neoplastic process, eventually leading to the development of pancreatic cancer^{49,50}. The same mutations occurring in normal colonic or ovarian epithelial cells lead to self-limiting hyperplastic or borderline lesions that do not progress to malignancy^{51–53}.

ii. Chronology: In contrast to the effects of *KRAS2* mutations in a normal colonic epithelial cell, a *KRAS2* gene mutation in the same cell type that has already acquired an *APC* mutation results in a clonal expansion that often progresses to cancer⁵⁴.

iii. Growth inhibition versus growth promotion: In many human and experimental cancers, *RAS* genes seem to function as oncogenes^{55,56}. But *RAS* genes can function as suppressor genes under other circumstances, inhibiting tumorigenesis after administration of carcinogens to mice^{57,58}.

These and similar observations on other cancer genes are consistent with the emerging general notion that signaling molecules play multiple roles at multiple times, even in the same cell type (for example, see ref. 59). However, the biochemical bases for such variations among cancer cells are almost entirely unknown. One could argue that oncogenes and tumor-suppressor genes are like electronic components whose effects depend on their placement within an electrical circuit. But no such argument can be easily used to explain the cell type specificity of stability gene defects. For example, MMR genes seem to have the identical, nonredundant function in every cell type on the planet: they limit the acquisition of specific types of mutations, such as those in homopolymer tracts^{60,61}. Yet inherited mismatch repair defects lead to tumors in the colon and endometrium but spare most other organs, including rapidly dividing, self-renewing tissues such as the small intestine and bone marrow^{62–64}.




Figure 7 PI3K pathway. Symbols as in Figure 1.

It is also notable in this context that many cancer genes affect different organs when mutated in mice than when mutated in humans⁸. An important practical implication of all this complexity is that it is risky to generalize the conclusions of experiments performed in any cell type or to infer the function of a gene in human cancer cells based on studies of the homologous genes in other organisms.

Major cast members or bit players? In addition to the genes that are mutated in a significant portion of cancers of a given type, such as those listed in Tables 1 and 2, there are many other genes that have been implicated in neoplasia but not shown to be mutated. These genes have been shown to be expressed at higher or lower levels than expected in normal cells^{65,66} and are often associated with ‘epigenetic’ changes—that is, covalent modifications of DNA or chromatin that are preserved as the cancer cells divide^{67,68}. Unlike genetic changes, epigenetic changes identical to those found in cancers are often found in normal cells at some stage of development.

The discovery of such genes is a growth industry now that high-throughput methods for evaluating the genes expressed in cancer cells have been developed^{65,66,69–74}. The information gained from these studies has proven extremely promising for the development of diagnostic assays, particularly for prognosis (for example, see ref. 75). But as has been shown for prostate-specific antigen (PSA), the utility of a gene for cancer diagnostics does not necessarily reflect a causative role in the process. It will therefore be essential to determine how genes discovered through expression-based approaches can be elevated from candidate status to culprit status in human neoplasia. When genes are sometimes inactivated by mutation and in other cases inactivated by epigenetic silencing, a




Figure 8 SMAD pathway. Symbols as in Figure 1.

cogent case can be made for their involvement in cancer. Relevant examples include the *VHL* and *CDKN2A* tumor-suppressor genes and the *hMLH1* stability gene⁶⁷. In the absence of mutational evidence, functional evidence obtained from studies *in vitro* or in nonhuman species *in vivo* must be used. But because of the issues with cell type, species and chronological specificity noted above, expression and functional studies cannot currently provide the ‘smoking guns’ that would definitively implicate the gene in human cancers^{76,77}. Telomerase provides a cogent example of the difficulties involved in such determinations. There is abundant evidence implicating telomerase in processes characteristic of neoplasia, such as immortalization^{45,78} and genetic instability^{42,79}, and a great deal is known about its biochemical properties^{80,81}. Even so, it is unclear at present whether the expression of telomerase in cancers is abnormal, driving the neoplastic process, or simply reflects the fact that cancers are derived from stem cells that normally express this enzyme.

Levels of gene expression are unreliable indicators of causation because disturbance of any network invariably leads to a multitude of such changes only peripherally related to the phenotype⁸². Without better ways to determine whether an unmutated but interesting candidate gene has a causal role in neoplasia, cancer researchers will likely be spending precious time working on genes only peripherally related to the disease they wish to study. One challenge for the future is therefore to develop new model systems. For liquid tumors, innovative systems have indeed been developed, such as those using immunodeficient mice reconstituted with human hematopoietic stem cells⁸³. Perhaps analogous humanized models can be developed to explore solid tumors in a more medically meaningful context than is currently possible^{84,85}.

What makes a box-office smash? Significant gains in cancer therapeutics have also been made in the last decade. Many of these advances have been incremental, but the increments add up. Thus more potent, less toxic derivatives of classic chemotherapeutic agents have been developed, dosing and combination treatments further optimized and side effects ameliorated⁸⁶. However, the




Figure 9 Overview of cancer gene pathways. The major pathways regulating cell birth and cell death are depicted as ovals color-coded to match Figs. 1–8. The schematics in Figs. 1–8 emphasize the genes that have been shown to be genetically altered in human tumors, though many other genes participate in these pathways. Additionally, some of the same genes appear in more than one pathway and there is substantial ‘cross-talk’ between pathways. Selected mediators of this cross-talk are indicated in the loops that connect the pathways. More detailed information about these pathways can be found in several comprehensive reviews (refs. 11,12,15,31,103–116).

Table 2 Genes that are mutated somatically but not inherited in mutant form

Gene ^a (synonym)	Somatic mutation type ^b	Cancers with mutant gene ^c	Pathway ^d
<i>CTNNB1</i> (β-catenin)	Activating codon change	Colon, liver, medulloblastomas	APC
<i>BCL2</i>	Translocation	Lymphomas	APOP
<i>TNFRSF6</i> (<i>FAS</i>)	Activating codon change	Lymphomas, testicular germ cell tumors	APOP
<i>BAX</i>	Inactivating codon change	Colon, stomach	APOP
<i>FBXW7</i> (<i>CDC4</i>)	Inactivating codon change	Colon, uterine, ovarian, breast	CIN
<i>GLI</i>	Amplification, translocation	Brain, sarcomas	GLI
<i>HPV E6</i>	HPV infection	Cervical	p53
<i>MDM2</i>	Amplification	Sarcomas	p53
<i>NOTCH1</i>	Translocation	Leukemias	p53
<i>AKT2</i>	Amplification	Ovarian, breast	PI3K
<i>FOXO1A, 3A</i>	Translocation	Rhabdomyosarcomas, leukemias	PI3K
<i>PI3KCA</i>	Activating codon change	Colon, stomach, brain, breast	PI3K
<i>CCND1</i> (cyclin D1)	Amplification, translocation	Leukemias, breast	RB
<i>HPV E7</i>	HPV infection	Cervical	RB
<i>TAL1</i>	Translocation	Leukemias	RB
<i>TFE3</i>	Translocation	Kidney, sarcomas	RB
<i>ABL1</i> (<i>ABL</i>)	Translocation	Chronic myelogenous leukemia	RTK
<i>ALK</i>	Translocation	Anaplastic large cell lymphoma	RTK
<i>BRAF</i>	Activating codon change	Melanoma, colorectal, thyroid	RTK
<i>EGFR</i>	Amplification, activating codon change	Glioblastomas, non-small cell lung cancers	RTK
<i>EPHB2</i>	Inactivating codon change	Prostate	RTK
<i>ERBB2</i>	Amplification	Breast, ovarian	RTK
<i>FES</i>	Activating codon change	Colon	RTK
<i>FGFR1–3</i>	Translocation	Lymphomas, gastric cancers, bladder cancers	RTK
<i>FLT3, 4</i>	Activating codon change	Leukemias, angiosarcomas	RTK
<i>JAK2</i>	Translocation	Leukemias	RTK
<i>KRAS2, N-RAS</i>	Activating codon change	Colorectal, pancreatic, non-small cell lung cancer	RTK
<i>NTRK1, 3</i>	Translocation, activating codon change	Thyroid, secretory breast, colon	RTK
<i>PDGFB</i>	Translocation	Dermatofibrosarcomas and fibroblastomas	RTK
<i>PDGFRB</i>	Translocation	Leukemias	RTK
<i>EWSR1</i>	Translocation	Ewing's sarcomas, lymphomas, leukemias	SMAD
<i>RUNX1</i>	Translocation	Leukemias	SMAD
<i>SMAD2</i>	Inactivating codon change	Colon, breast	SMAD
<i>TGFBR1, TGFBR2</i>	Inactivating codon change	Colon, stomach, ovarian	SMAD
<i>BCL6</i>	Translocation	Lymphomas	?
<i>EV11</i>	Translocation	Leukemias	?
<i>HMGA2</i>	Translocation	Lipomas	?
<i>HOXA9, 11, 13; HOXC13,</i> <i>HOXD11, 13; HOX11, HOX11L2</i>	Translocation	Leukemias	?
<i>MAP2K4</i> (<i>MKK4</i>)	Inactivating codon change	Pancreas, breast, colon	?
<i>MLL</i>	Translocation, activating codon change	Leukemias	?
<i>MYC, MYCN, MYCL1</i>	Amplification	Lymphomas, neuroblastomas, small cell lung cancers	?
<i>PTNP1, 11</i>	Activating codon change	Leukemias, colon	?
<i>RARA</i>	Translocation	Promyelocytic leukemia	?
<i>SS18</i>	Translocation	Synovial sarcomas	?

^aRepresentative genes of all the major pathways and cancer types are listed. For a complete list, see ref. 76. Approved gene symbols are provided for each entry, with alternative names in parentheses. ^bActivating codon change, intragenic mutation altering one or a small number of base pairs that activates the gene product, indicating that it is an oncogene; inactivating codon change, any mutation (point mutation, small or large deletion, etc.) that inactivates the gene product, indicating that the gene is a tumor suppressor. Amplifications and translocations generally affect oncogenes, though occasional translocations disrupt a gene rather than activate it (such as has been suggested to occur with *RUNX1*; ref. 118). ^cOnly representative types of cancers are listed when a gene is mutated in many tumor types. Specific types of leukemias and lymphomas are listed only if a given gene is predominantly mutated in a specific subtype. ^dIn many cases, the gene has been implicated in several pathways. The single pathway that is listed for each gene represents a 'best guess' (when one can be made) and for the reasons noted in the text and in the legend to **Figure 9**, should not be regarded as conclusive. APOP, apoptotic pathway; RTK, receptor tyrosine kinase pathway (see **Figure 9**).

most dramatic therapeutic advances have come from agents targeted against proteins encoded by genes that are mutated in cancers. These include trastuzumab (Herceptin), an antibody against the product of the *ERBB2* gene amplified in some breast cancers⁸⁷; imatinib (Gleevec), an inhibitor of tyrosine kinases altered in chronic myelogenous leukemia (CML)⁸⁸ and gastrointestinal stromal tumors (GISTs)⁸⁹; and gefitinib (Iressa), an EGFR

kinase inhibitor used to treat lung cancers⁹⁰. Though none of these advances have resulted in cures of many patients with advanced disease, they can substantially improve and prolong lives. One important lesson from the use of these agents is that mutations are more reliable indicators of a good target than is abnormal expression. For example, virtually all GISTs abnormally express high levels of c-kit, but only those tumors with intragenic mutations of *KIT*

respond to therapy with imatinib⁹¹. Similarly, gefitinib is efficacious only in lung cancers in which the kinase domain of *EGFR* is mutated but not in lung cancers in which *EGFR* is simply overexpressed^{92,93}.

Interestingly, we still don't know the basis for the selectivity of conventional chemotherapeutic agents. Why cancer cells are more sensitive to antimetabolites, alkylating agents, DNA intercalators and topoisomerase inhibitors than normal replicating cells is enigmatic. The explosion in research on apoptosis in the 1990s has suggested that some tumor cells are less, rather than more, likely to undergo apoptosis after noxious stimuli^{94,95}. Differential apoptotic proclivities therefore don't readily explain therapeutic specificities. This is consistent with studies showing that many drugs used in the clinic to treat solid tumors do not function through the induction of apoptosis at the doses achieved in patients⁹⁶. Even with the newer, targeted agents, the basis for selectivity is not generally clear. The ERBB2 and ABL tyrosine kinases, for example, are expressed in many normal cell types; why should targeting these kinases lead to the demise of cancer cells but not normal cells? Concepts such as 'cellular addiction' offer an imaginative answer to this question, but the molecular basis for the postulated addiction is unknown⁹⁷. Similar arguments could be made for 'untargeted' drugs like Taxol; are cancer cells addicted to microtubules? It is interesting that so much work has been done on how cells become resistant to conventional chemotherapeutic agents and so little on why they are sensitive to begin with. Indeed, some studies indicate that it is not the neoplastic cells, but rather the tumor vasculature, that is the actual target of many conventional chemotherapeutic agents²⁷. Perhaps if we understood the bases for the responsiveness of cancers to these drugs, we could devise better agents to exploit them.

The sequel. There are at least three major challenges that will occupy most cancer researchers' time over the next 10 years. The first is the discovery of new genes that have a causal role in neoplasia, particularly those that initiate and conclude the process. The second is the delineation of the pathways through which these genes act and the basis for the varying actions in specific cell types. The third is the development of new ways to exploit this knowledge for the benefit of patients.

The first two of these challenges are likely to proceed apace. Technologies for gene discovery are rapidly advancing, the human genome will soon be in finished form, and previous analyses of other genes provide well-traveled road maps for investigators to follow once new genes are in hand. In fact, there are likely to be too many genes, too many interacting proteins and too many potential functions to consider, rather than too few. As noted above, cancer biology has not kept up with cancer molecular genetics and new biological systems are needed to separate wheat from chaff.

The third challenge, involving practical benefits to patients, will be much more difficult to meet. There are hardly any road maps to follow. Eradicating hundreds of billions of cancer cells from a human with metastatic disease is a daunting task. Each of these cells has multiple genetic abnormalities and is capable of rapidly evolving variants to combat any therapeutic onslaught^{98,99}. On the other hand, many cancers respond to the somewhat crude weapons that have been developed to date despite their multiple genetic abnormalities. And now that cancer is no longer a 'black box', and is known to be the result of alterations in a limited number of pathways that can in principle be targeted by new generations of drugs, cautious optimism is appropriate.

But will the development of new, more specific therapeutic agents be the best way to minimize cancer morbidity and mortality in the long-term? Most of the increased longevity that Western societies enjoy today has come through better prevention rather than better treatment. It is clear that certain major forms of cancer can be prevented by limiting exposures to carcinogens (such as sunlight and cigarette smoke in skin and lung cancers, respectively). Other forms of cancer can be detected early, thereby limiting morbidity and mortality (for example, tumors of the breast, cervix, prostate and colon). Moreover, it requires 30–40 years for a typical epithelial cell to accumulate the multiple genetic alterations required to progress to metastatic disease¹⁰⁰. This provides a huge window of opportunity to detect tumors at a stage when they are still curable by conventional surgical methods. Perhaps knowledge of the genes altered in cancer will provide the basis for new generations of diagnostic tests, employing target-specific imaging or analysis of body fluid samples *ex vivo* that will make early detection approaches feasible^{101,102}. Though less dramatic than cures, prevention and early detection are perhaps the most promising and feasible means to reduce cancer deaths by the time that *Nature Medicine* celebrates its 20th anniversary.

ACKNOWLEDGMENTS

The authors thank many colleagues for their critical reading of the manuscript and S. Mousset for sharing results on EphB2 before publication. Work in the authors' laboratories has been supported by the Ludwig Trust, the National Colorectal Cancer Research Alliance, the Miracle Foundation, the Clayton Fund and US National Institutes of Health grants CA43460, CA57345 and CA62924.

COMPETING INTERESTS STATEMENT

The authors declare competing financial interests (see the *Nature Medicine* website for details).

This material is part of Nature Medicine's 10 year anniversary series. For more content related to these special focus issues, please see http://www.nature.com/nm/special_focus/anniversary/index.html

Published online at <http://www.nature.com/naturemedicine/>

- Davies, H. *et al.* Mutations of the BRAF gene in human cancer. *Nature* **417**, 949–954 (2002).
- Wan, P.T. *et al.* Mechanism of activation of the RAF-ERK signaling pathway by oncogenic mutations of B-RAF. *Cell* **116**, 855–867 (2004).
- Santarosa, M. & Ashworth, A. Haploinsufficiency for tumour suppressor genes: when you don't need to go all the way. *Biochim. Biophys. Acta* **1654**, 105–122 (2004).
- Knudson, A.G. Cancer genetics. *Am. J. Med. Genet.* **111**, 96–102 (2002).
- Friedberg, E.C. DNA damage and repair. *Nature* **421**, 436–440 (2003).
- Nowell, P.C. Tumor progression: a brief historical perspective. *Semin. Cancer Biol.* **12**, 261–266 (2002).
- Maley, C.C. *et al.* Selectively advantageous mutations and hitchhikers in neoplasms: p16 lesions are selected in Barrett's esophagus. *Cancer Res.* **64**, 3414–3427 (2004).
- Van Dyke, T. & Jacks, T. Cancer modeling in the modern era: progress and challenges. *Cell* **108**, 135–144 (2002).
- Horvitz, H.R. Worms, life, and death. *ChemBiochem* **4**, 697–711 (2003).
- Sherr, C.J. Cancer cell cycles revisited. *Cancer Res.* **60**, 3689–3695 (2000).
- Ortega, S., Malumbres, M. & Barbacid, M. Cyclin D-dependent kinases, INK4 inhibitors and cancer. *Biochim. Biophys. Acta* **1602**, 73–87 (2002).
- Classon, M. & Harlow, E. The retinoblastoma tumour suppressor in development and cancer. *Nat. Rev. Cancer* **2**, 910–917 (2002).
- Ichimura, K. *et al.* Deregulation of the p14ARF/MDM2/p53 pathway is a prerequisite for human astrocytic gliomas with G1-S transition control gene abnormalities. *Cancer Res.* **60**, 417–424 (2000).
- Vogelstein, B., Lane, D. & Levine, A.J. Surfing the p53 network. *Nature* **408**, 307–310 (2000).
- Oren, M. Decision making by p53: life, death and cancer. *Cell Death Differ.* **10**, 431–442 (2003).
- Prives, C. & Hall, P.A. The p53 pathway. *J. Pathol.* **187**, 112–126 (1999).
- Klein, G. Perspectives in studies of human tumor viruses. *Front. Biosci.* **7**, d268–d274 (2002).
- Munger, K. & Howley, P.M. Human papillomavirus immortalization and transformation functions. *Virus Res.* **89**, 213–228 (2002).
- zur Hausen, H. Oncogenic DNA viruses. *Oncogene* **20**, 7820–7823 (2001).

20. Hunter, T. Signaling—2000 and beyond. *Cell* **100**, 113–127 (2000).
21. Komarova, N.L., Sengupta, A. & Nowak, M.A. Mutation-selection networks of cancer initiation: tumor suppressor genes and chromosomal instability. *J. Theor. Biol.* **223**, 433–450 (2003).
22. Rowley, J.D. The critical role of chromosome translocations in human leukemias. *Annu. Rev. Genet.* **32**, 495–519 (1998).
23. Mitelman, F. Recurrent chromosome aberrations in cancer. *Mutat. Res.* **462**, 247–253 (2000).
24. Verheul, H.M., Voest, E.E. & Schlingemann, R.O. Are tumours angiogenesis-dependent? *J. Pathol.* **202**, 5–13 (2004).
25. Tlsty, T.D. & Hein, P.W. Know thy neighbor: stromal cells can contribute oncogenic signals. *Curr. Opin. Genet. Dev.* **11**, 54–59 (2001).
26. Fata, J.E., Werb, Z. & Bissell, M.J. Regulation of mammary gland branching morphogenesis by the extracellular matrix and its remodeling enzymes. *Breast Cancer Res.* **6**, 1–11 (2004).
27. Kerbel, R. & Folkman, J. Clinical translation of angiogenesis inhibitors. *Nat. Rev. Cancer* **2**, 727–739 (2002).
28. Folkman, J. Role of angiogenesis in tumor growth and metastasis. *Semin. Oncol.* **29**, 15–18 (2002).
29. Ferrara, N., Hillan, K.J., Gerber, H.P. & Novotny, W. Discovery and development of bevacizumab, an anti-VEGF antibody for treating cancer. *Nat. Rev. Drug Discov.* **3**, 391–400 (2004).
30. Kondo, K., Kiko, J., Nakamura, E., Lechpammer, M. & Kaelin, W.G. Jr. Inhibition of HIF is necessary for tumor suppression by the von Hippel-Lindau protein. *Cancer Cell* **1**, 237–246 (2002).
31. Semenza, G.L. Targeting HIF-1 for cancer therapy. *Nat. Rev. Cancer* **3**, 721–732 (2003).
32. Strausberg, R.L., Simpson, A.J. & Wooster, R. Sequence-based cancer genomics: progress, lessons and opportunities. *Nat. Rev. Genet.* **4**, 409–418 (2003).
33. Loeb, L.A., Loeb, K.R. & Anderson, J.P. Multiple mutations and cancer. *Proc. Natl. Acad. Sci. USA* **100**, 776–781 (2003).
34. Rajagopalan, H., Nowak, M.A., Vogelstein, B. & Lengauer, C. The significance of unstable chromosomes in colorectal cancer. *Nat. Rev. Cancer* **3**, 695–701 (2003).
35. Sieber, O.M., Heinemann, K. & Tomlinson, I.P. Genomic instability—the engine of tumorigenesis? *Nat. Rev. Cancer* **3**, 701–708 (2003).
36. Wang, T.L. *et al.* Prevalence of somatic alterations in the colorectal cancer cell genome. *Proc. Natl. Acad. Sci. USA* **99**, 3076–3080 (2002).
37. Lengauer, C., Kinzler, K.W. & Vogelstein, B. Genetic instabilities in human cancers. *Nature* **396**, 643–649 (1998).
38. Duesberg, P. & Li, R. Multistep carcinogenesis: a chain reaction of aneuploidizations. *Cell Cycle* **2**, 202–210 (2003).
39. Albertson, D.G. & Pinkel, D. Genomic microarrays in human genetic disease and cancer. *Hum. Mol. Genet.* **12** (spec. no. 2), R145–R152 (2003).
40. Shiloh, Y. & Kastan, M.B. ATM: genome stability, neuronal development, and cancer cross paths. *Adv. Cancer Res.* **83**, 209–254 (2001).
41. Scully, R. & Livingston, D.M. In search of the tumour-suppressor functions of BRCA1 and BRCA2. *Nature* **408**, 429–432 (2000).
42. Maser, R.S. & DePinho, R.A. Connecting chromosomes, crisis, and cancer. *Science* **297**, 565–569 (2002).
43. Pihan, G. & Doxsey, S.J. Mutations and aneuploidy: co-conspirators in cancer? *Cancer Cell* **4**, 89–94 (2003).
44. Rajagopalan, H. *et al.* Inactivation of hCDC4 can cause chromosomal instability. *Nature* **428**, 77–81 (2004).
45. Shay, J.W. & Roninson, I.B. Hallmarks of senescence in carcinogenesis and cancer therapy. *Oncogene* **23**, 2919–2933 (2004).
46. Chambers, A.F., Groom, A.C. & MacDonald, I.C. Dissemination and growth of cancer cells in metastatic sites. *Nat. Rev. Cancer* **2**, 563–572 (2002).
47. Fidler, I.J. Critical determinants of metastasis. *Semin. Cancer Biol.* **12**, 89–96 (2002).
48. Hunter, K.W. Host genetics and tumour metastasis. *Br. J. Cancer* **90**, 752–755 (2004).
49. Hruban, R.H., Goggins, M., Parsons, J. & Kern, S.E. Progression model for pancreatic cancer. *Clin. Cancer Res.* **6**, 2969–2972 (2000).
50. Aguirre, A.J. *et al.* Activated Kras and Ink4a/Arf deficiency cooperate to produce metastatic pancreatic ductal adenocarcinoma. *Genes Dev.* **17**, 3112–3126 (2003).
51. Jen, J. *et al.* Molecular determinants of dysplasia in colorectal lesions. *Cancer Res.* **54**, 5523–5526 (1994).
52. Pretlow, T.P. Aberrant crypt foci and K-ras mutations: earliest recognized players or innocent bystanders in colon carcinogenesis? *Gastroenterology* **108**, 600–603 (1995).
53. Sieben, N.L. *et al.* In ovarian neoplasms, BRAF, but not KRAS, mutations are restricted to low-grade serous tumours. *J. Pathol.* **202**, 336–340 (2004).
54. Kinzler, K.W. & Vogelstein, B. Colorectal Tumors. In *The Genetic Basis of Human Cancer* (eds. Vogelstein, B. & Kinzler, K.W.) 565–587 (McGraw-Hill, New York, 1998).
55. Barbacid, M. ras genes. *Annu. Rev. Biochem.* **56**, 779–827 (1987).
56. Bos, J.L. ras oncogenes in human cancer: a review. *Cancer Res.* **49**, 4682–4689 (1989).
57. Zhang, Z. *et al.* Wildtype Kras2 can inhibit lung carcinogenesis in mice. *Nat. Genet.* **29**, 25–33 (2001).
58. Diaz, R. *et al.* The N-ras proto-oncogene can suppress the malignant phenotype in the presence or absence of its oncogene. *Cancer Res.* **62**, 4514–4518 (2002).
59. Bronner-Fraser, M. Development. Making sense of the sensory lineage. *Science* **303**, 966–968 (2004).
60. Jiricny, J. Eukaryotic mismatch repair: an update. *Mutat. Res.* **409**, 107–121 (1998).
61. Fishel, R. & Wilson, T. MutS homologs in mammalian cells. *Curr. Opin. Genet. Dev.* **7**, 105–113 (1997).
62. Lynch, H.T. & de la Chapelle, A. Hereditary colorectal cancer. *N. Engl. J. Med.* **348**, 919–932 (2003).
63. Yamamoto, H., Imai, K. & Perucho, M. Gastrointestinal cancer of the microsatellite mutator phenotype pathway. *J. Gastroenterol.* **37**, 153–163 (2002).
64. Honchel, R., Halling, K.C. & Thibodeau, S.N. Genomic instability in neoplasia. *Semin. Cell Biol.* **6**, 45–52 (1995).
65. Brown, P.O. & Botstein, D. Exploring the new world of the genome with DNA microarrays. *Nat. Genet.* **21**, 33–37 (1999).
66. Polyak, K. & Riggins, G.J. Gene discovery using the serial analysis of gene expression technique: implications for cancer research. *J. Clin. Oncol.* **19**, 2948–2958 (2001).
67. Jones, P.A. & Baylin, S.B. The fundamental role of epigenetic events in cancer. *Nat. Rev. Genet.* **3**, 415–428 (2002).
68. Feinberg, A.P. & Tycko, B. The history of cancer epigenetics. *Nat. Rev. Cancer* **4**, 143–153 (2004).
69. Collins, F.S., Green, E.D., Guttmacher, A.E. & Guyer, M.S. A vision for the future of genomics research. *Nature* **422**, 835–847 (2003).
70. Schadt, E.E., Monks, S.A. & Friend, S.H. A new paradigm for drug discovery: integrating clinical, genetic, genomic and molecular phenotype data to identify drug targets. *Biochem. Soc. Trans.* **31**, 437–443 (2003).
71. Paddison, P.J. *et al.* A resource for large-scale RNA-interference-based screens in mammals. *Nature* **428**, 427–431 (2004).
72. Berns, K. *et al.* A large-scale RNAi screen in human cells identifies new components of the p53 pathway. *Nature* **428**, 431–437 (2004).
73. Rosenblatt, K.P. *et al.* Serum proteomics in cancer diagnosis and management. *Annu. Rev. Med.* **55**, 97–112 (2004).
74. Luo, J., Isaacs, W.B., Trent, J.M. & Duggan, D.J. Looking beyond morphology: cancer gene expression profiling using DNA microarrays. *Cancer Invest.* **21**, 937–949 (2003).
75. Ma, X.J. *et al.* A two-gene expression ratio predicts clinical outcome in breast cancer patients treated with tamoxifen. *Cancer Cell* **5**, 607–616 (2004).
76. Futreal, P.A. *et al.* A census of human cancer genes. *Nat. Rev. Cancer* **4**, 177–183 (2004).
77. Masayeva, B.G. *et al.* Gene expression alterations over large chromosomal regions in cancers include multiple genes unrelated to malignant progression. *Proc. Natl. Acad. Sci. USA* **101**, 8715–8720 (2004).
78. Stewart, S.A. & Weinberg, R.A. Senescence: does it all happen at the ends? *Oncogene* **21**, 627–630 (2002).
79. Feldser, D.M., Hackett, J.A. & Greider, C.W. Telomere dysfunction and the initiation of genome instability. *Nat. Rev. Cancer* **3**, 623–627 (2003).
80. Chan, S.R. & Blackburn, E.H. Telomeres and telomerase. *Phil. Trans. R. Soc. Lond. B* **359**, 109–121 (2004).
81. Cech, T.R. Beginning to understand the end of the chromosome. *Cell* **116**, 273–279 (2004).
82. Miklos, G.L. & Maleszka, R. Microarray reality checks in the context of a complex disease. *Nat. Biotechnol.* **22**, 615–621 (2004).
83. Hope, K.J., Jin, L. & Dick, J.E. Human acute myeloid leukemia stem cells. *Arch. Med. Res.* **34**, 507–514 (2003).
84. Berking, C. & Herlyn, M. Human skin reconstruct models: a new application for studies of melanocyte and melanoma biology. *Histol. Histopathol.* **16**, 669–674 (2001).
85. Kuperwasser, C. *et al.* Reconstruction of functionally normal and malignant human breast tissues in mice. *Proc. Natl. Acad. Sci. USA* **101**, 4966–4971 (2004).
86. Frei, E.I. & Eder, J.P. Principles of dose, schedule, and combination Therapy. In *Cancer Medicine* (eds. Kufe, D.W. *et al.*) 669–677 (B.C. Decker, Inc., Hamilton, Ontario, 2003).
87. Pegram, M.D., Konecny, G. & Slamon, D.J. The molecular and cellular biology of HER2/neu gene amplification/overexpression and the clinical development of herceptin (trastuzumab) therapy for breast cancer. *Cancer Treat. Res.* **103**, 57–75 (2000).
88. Druker, B.J. *et al.* Chronic myelogenous leukemia. In *Hematology 2001 (American Society of Hematology Education Program)* 87–112 (American Society of Hematology, 2001).
89. Mechttersheimer, G. *et al.* Gastrointestinal stromal tumours and their response to treatment with the tyrosine kinase inhibitor imatinib. *Virchows Arch.* **444**, 108–118 (2004).
90. Langer, C.J. Emerging role of epidermal growth factor receptor inhibition in therapy for advanced malignancy: focus on NSCLC. *Int. J. Radiat. Oncol. Biol. Phys.* **58**, 991–1002 (2004).
91. Duensing, A., Heinrich, M.C., Fletcher, C.D. & Fletcher, J.A. Biology of gastrointestinal stromal tumors: KIT mutations and beyond. *Cancer Invest.* **22**, 106–116 (2004).
92. Paez, J.G. *et al.* EGFR mutations in lung cancer: correlation with clinical response to gefitinib therapy. *Science* **304**, 1497–1500 (2004).

93. Lynch, T.J. *et al.* Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. *N. Engl. J. Med.* **350**, 2129–2139 (2004).
94. Schmitt, C.A. & Lowe, S.W. Apoptosis and therapy. *J. Pathol.* **187**, 127–137 (1999).
95. Danial, N.N. & Korsmeyer, S.J. Cell death: critical control points. *Cell* **116**, 205–219 (2004).
96. Brown, J.M. & Wouters, B.G. Apoptosis: mediator or mode of cell killing by anticancer agents? *Drug Resist. Updat.* **4**, 135–136 (2001).
97. Weinstein, I.B. *et al.* Disorders in cell circuitry associated with multistage carcinogenesis: exploitable targets for cancer prevention and therapy. *Clin. Cancer Res.* **3**, 2696–2702 (1997).
98. Nygren, P. & Larsson, R. Overview of the clinical efficacy of investigational anticancer drugs. *J. Intern. Med.* **253**, 46–75 (2003).
99. Shih, L.Y. *et al.* Heterogeneous patterns of FLT3 Asp(835) mutations in relapsed de novo acute myeloid leukemia: a comparative analysis of 120 paired diagnostic and relapse bone marrow samples. *Clin. Cancer Res.* **10**, 1326–1332 (2004).
100. Kinzler, K.W. & Vogelstein, B. Lessons from hereditary colon cancer. *Cell* **87**, 159–170 (1996).
101. Weissleder, R. & Ntziachristos, V. Shedding light onto live molecular targets. *Nat. Med.* **9**, 123–128 (2003).
102. Sidransky, D. Emerging molecular markers of cancer. *Nat. Rev. Cancer* **2**, 210–219 (2002).
103. Gschwind, A., Fischer, O.M. & Ullrich, A. The discovery of receptor tyrosine kinases: targets for cancer therapy. *Nat. Rev. Cancer* **4**, 361–370 (2004).
104. Downward, J. Targeting RAS signalling pathways in cancer therapy. *Nat. Rev. Cancer* **3**, 11–22 (2003).
105. Malumbres, M. & Barbacid, M. To cycle or not to cycle: a critical decision in cancer. *Nat. Rev. Cancer* **1**, 222–231 (2001).
106. Giles, R.H., van Es, J.H. & Clevers, H. Caught up in a Wnt storm: Wnt signaling in cancer. *Biochim. Biophys. Acta* **1653**, 1–24 (2003).
107. Cantley, L.C. The phosphoinositide 3-kinase pathway. *Science* **296**, 1655–1657 (2002).
108. Shi, Y. & Massague, J. Mechanisms of TGF- β signaling from cell membrane to the nucleus. *Cell* **113**, 685–700 (2003).
109. Ruiz i Altaba, A., Stecca, B. & Sanchez, P. Hedgehog–Gli signaling in brain tumors: stem cells and paradevelopmental programs in cancer. *Cancer Lett.* **204**, 145–157 (2004).
110. Adams, J.M. Ways of dying: multiple pathways to apoptosis. *Genes Dev.* **17**, 2481–2495 (2003).
111. Blagosklonny, M.V. & Pardee, A.B. The restriction point of the cell cycle. *Cell Cycle* **1**, 103–110 (2002).
112. Plas, D.R. & Thompson, C.B. Cell metabolism in the regulation of programmed cell death. *Trends Endocrinol. Metab.* **13**, 75–78 (2002).
113. Green, D.R. & Evan, G.I. A matter of life and death. *Cancer Cell* **1**, 19–30 (2002).
114. Eng, C., Kiuru, M., Fernandez, M.J. & Aaltonen, L.A. A role for mitochondrial enzymes in inherited neoplasia and beyond. *Nat. Rev. Cancer* **3**, 193–202 (2003).
115. Lum, L. & Beachy, P.A. The Hedgehog response network: sensors, switches, and routers. *Science* **304**, 1755–1759 (2004).
116. Brivanlou, A.H. & Darnell, J.E. Jr. Signal transduction and the control of gene expression. *Science* **295**, 813–818 (2002).
117. Vogelstein, B. & Kinzler, K.W. *The Genetic Basis of Human Cancer* (McGraw-Hill, Toronto, 2002).
118. Cameron, E.R. & Neil, J.C. The Runx genes: lineage-specific oncogenes and tumor suppressors. *Oncogene* **23**, 4308–4314 (2004).



Cancer Genome Landscapes

Bert Vogelstein *et al.*

Science **339**, 1546 (2013);

DOI: 10.1126/science.1235122

This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

The following resources related to this article are available online at www.sciencemag.org (this information is current as of May 22, 2013):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/339/6127/1546.full.html>

Supporting Online Material can be found at:

<http://www.sciencemag.org/content/suppl/2013/03/27/339.6127.1546.DC1.html>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/content/339/6127/1546.full.html#related>

This article **cites 152 articles**, 57 of which can be accessed free:

<http://www.sciencemag.org/content/339/6127/1546.full.html#ref-list-1>

This article appears in the following **subject collections**:

Medicine, Diseases

<http://www.sciencemag.org/cgi/collection/medicine>

REVIEW

Cancer Genome Landscapes

Bert Vogelstein, Nickolas Papadopoulos, Victor E. Velculescu, Shibin Zhou, Luis A. Diaz Jr., Kenneth W. Kinzler*

Over the past decade, comprehensive sequencing efforts have revealed the genomic landscapes of common forms of human cancer. For most cancer types, this landscape consists of a small number of “mountains” (genes altered in a high percentage of tumors) and a much larger number of “hills” (genes altered infrequently). To date, these studies have revealed ~140 genes that, when altered by intragenic mutations, can promote or “drive” tumorigenesis. A typical tumor contains two to eight of these “driver gene” mutations; the remaining mutations are passengers that confer no selective growth advantage. Driver genes can be classified into 12 signaling pathways that regulate three core cellular processes: cell fate, cell survival, and genome maintenance. A better understanding of these pathways is one of the most pressing needs in basic cancer research. Even now, however, our knowledge of cancer genomes is sufficient to guide the development of more effective approaches for reducing cancer morbidity and mortality.

Ten years ago, the idea that all of the genes altered in cancer could be identified at base-pair resolution would have seemed like science fiction. Today, such genome-wide analysis, through sequencing of the exome (see Box 1, Glossary, for definitions of terms used in this Review) or of the whole genome, is routine.

The prototypical exomic studies of cancer evaluated ~20 tumors at a cost of >\$100,000 per case (1–3). Today, the cost of this sequencing has been reduced 100-fold, and studies reporting the sequencing of more than 100 tumors of a given type are the norm (table S1A). Although vast amounts of data can now be readily obtained, deciphering this information in meaningful terms is still challenging. Here, we review what has been learned about cancer genomes from these sequencing studies—and, more importantly, what this information has taught us about cancer biology and future cancer management strategies.

How Many Genes Are Subtly Mutated in a Typical Human Cancer?

In common solid tumors such as those derived from the colon, breast, brain, or pancreas, an average of 33 to 66 genes display subtle somatic mutations that would be expected to alter their protein products (Fig. 1A). About 95% of these mutations are single-base substitutions (such as C>G), whereas the remainder are deletions or insertions of one or a few bases (such as CTT>CT) (table S1B). Of the base substitutions, 90.7% result in missense changes, 7.6% result in nonsense changes, and 1.7% result in alterations of splice sites or untranslated regions immediately adjacent to the start and stop codons (table S1B).

The Ludwig Center and The Howard Hughes Medical Institute at Johns Hopkins Kimmel Cancer Center, Baltimore, MD 21287, USA.

*Corresponding author. E-mail: kinzkle@jhmi.edu

Certain tumor types display many more or many fewer mutations than average (Fig. 1B). Notable among these outliers are melanomas and lung tumors, which contain ~200 nonsynonymous mutations per tumor (table S1C). These larger numbers reflect the involvement of potent mutagens (ultraviolet light and cigarette smoke, respectively) in the pathogenesis of these tumor types. Accordingly, lung cancers from smokers have 10 times as many somatic mutations as those from nonsmokers (4). Tumors with defects in DNA repair form another group of outliers (5). For example, tumors with mismatch repair defects can harbor thousands of mutations (Fig. 1B), even more than lung tumors or melanomas. Recent studies have shown that high numbers of mutations are also found in tumors with genetic alterations of the proofreading domain of DNA polymerases POLE or POLD1 (6, 7). At the other end of the spectrum, pediatric tumors and leukemias harbor far fewer point mutations: on average, 9.6 per tumor (table S1C). The basis for this observation is considered below.

Mutation Timing

When do these mutations occur? Tumors evolve from benign to malignant lesions by acquiring a series of mutations over time, a process that has been particularly well studied in colorectal tumors (8, 9). The first, or “gatekeeping,” mutation provides a selective growth advantage to a normal epithelial cell, allowing it to outgrow the cells that surround it and become a microscopic clone (Fig. 2). Gatekeeping mutations in the colon most often occur in the *APC* gene (10). The small adenoma that results from this mutation grows slowly, but a second mutation in another gene, such as *KRAS*, unleashes a second round of clonal growth that allows an expansion of cell number (9). The cells with only the *APC* mutation may persist, but their cell numbers are small compared with the cells that

have mutations in both genes. This process of mutation followed by clonal expansion continues, with mutations in genes such as *PIK3CA*, *SMAD4*, and *TP53*, eventually generating a malignant tumor that can invade through the underlying basement membrane and metastasize to lymph nodes and distant organs such as the liver (11). The mutations that confer a selective growth advantage to the tumor cell are called “driver” mutations. It has been estimated (12) that each driver mutation provides only a small selective growth advantage to the cell, on the order of a 0.4% increase in the difference between cell birth and cell death. Over many years, however, this slight increase, compounded once or twice per week, can result in a large mass, containing billions of cells.

The number of mutations in certain tumors of self-renewing tissues is directly correlated with age (13). When evaluated through linear regression, this correlation implies that more than half of the somatic mutations identified in these tumors occur during the preneoplastic phase; that is, during the growth of normal cells that continuously replenish gastrointestinal and genitourinary epithelium and other tissues. All of these pre-neoplastic mutations are “passenger” mutations that have no effect on the neoplastic process. This result explains why a colorectal tumor in a 90-year-old patient has nearly twice as many mutations as a morphologically identical colorectal tumor in a 45-year-old patient. This finding also partly explains why advanced brain tumors (glioblastomas) and pancreatic cancers (pancreatic ductal adenocarcinomas) have fewer mutations than colorectal tumors; glial cells of the brain and epithelial cells of the pancreatic ducts do not replicate, unlike the epithelial cells lining the crypts of the colon. Therefore, the gatekeeping mutation in a pancreatic or brain cancer is predicted to occur in a precursor cell that contains many fewer mutations than are present in a colorectal precursor cell. This line of reasoning also helps to explain why pediatric cancers have fewer mutations than adult tumors. Pediatric cancers often occur in non-self-renewing tissues, and those that arise in renewing tissues (such as leukemias) originate from precursor cells that have not renewed themselves as often as in adults. In addition, pediatric tumors, as well as adult leukemias and lymphomas, may require fewer rounds of clonal expansion than adult solid tumors (8, 14). Genome sequencing studies of leukemia patients support the idea that mutations occur as random events in normal precursor cells before these cells acquire an initiating mutation (15).

When during tumorigenesis do the remaining somatic mutations occur? Because mutations in tumors occur at predictable and calculable rates (see below), the number of somatic mutations in tumors provides a clock, much like the clock used in evolutionary biology to determine species

divergence time. The number of mutations has been measured in tumors representing progressive stages of colorectal and pancreatic cancers (11, 16). Applying the evolutionary clock model to these data leads to two unambiguous conclusions: First, it takes decades to develop a full-blown, metastatic cancer. Second, virtually all of the mutations in metastatic lesions were already present in a large number of cells in the primary tumors.

The timing of mutations is relevant to our understanding of metastasis, which is responsible for the death of most patients with cancer. The primary tumor can be surgically removed, but the residual metastatic lesions—often undetectable and widespread—remain and eventually enlarge, compromising the function of the lungs, liver, or other organs. From a genetics perspective, it would seem that there must be mutations that convert a primary cancer to a metastatic one, just as there are mutations that convert a normal cell to a benign tumor, or a benign tumor to a malignant one (Fig. 2). Despite intensive effort, however, consistent genetic alterations that distinguish cancers that metastasize from cancers that have not yet metastasized remain to be identified.

One potential explanation invokes mutations or epigenetic changes that are difficult to identify with current technologies (see section on “dark matter” below). Another explanation is that metastatic lesions have not yet been studied in sufficient detail to identify these genetic alterations, particularly if the mutations are heterogeneous in nature. But another possible explanation is that there are no metastasis genes. A malignant primary tumor can take many years to metastasize, but this process is, in principle, explicable by stochastic processes alone (17, 18). Advanced tumors release millions of cells into the circulation each day, but these cells have short half-lives, and only a minuscule fraction establish metastatic lesions (19). Conceivably, these circulating cells may, in a nondeterministic manner, infrequently and randomly lodge in a capillary bed in an organ that provides a favorable microenvironment for growth. The bigger the primary tumor mass, the more likely that this process will occur. In this scenario, the continual evolution of the primary tumor would reflect local selective advantages rather than future selective advantages. The idea that growth at metastatic sites is not dependent on additional genetic alterations is also supported by recent results showing that even normal cells, when placed in suitable environments such as lymph nodes, can grow into organoids, complete with a functioning vasculature (20).

Other Types of Genetic Alterations in Tumors

Though the rate of point mutations in tumors is similar to that of normal cells, the rate of chromosomal changes in cancer is elevated (21). Therefore, most solid tumors display widespread changes in chromosome number (aneuploidy), as well as deletions, inversions, translocations,




Fig. 1. Number of somatic mutations in representative human cancers, detected by genome-wide sequencing studies. (A) The genomes of a diverse group of adult (right) and pediatric (left) cancers have been analyzed. Numbers in parentheses indicate the median number of nonsynonymous mutations per tumor. **(B)** The median number of nonsynonymous mutations per tumor in a variety of tumor types. Horizontal bars indicate the 25 and 75% quartiles. MSI, microsatellite instability; SCLC, small cell lung cancers; NSCLC, non-small cell lung cancers; ESCC, esophageal squamous cell carcinomas; MSS, microsatellite stable; EAC, esophageal adenocarcinomas. The published data on which this figure is based are provided in table S1C.

CREDIT: FIG. 1A, I.E. COOK




Fig. 2. Genetic alterations and the progression of colorectal cancer. The major signaling pathways that drive tumorigenesis are shown at the transitions between each tumor stage. One of several driver genes that encode compo-

nents of these pathways can be altered in any individual tumor. Patient age indicates the time intervals during which the driver genes are usually mutated. Note that this model may not apply to all tumor types. TGF- β , transforming growth factor- β .

and other genetic abnormalities. When a large part of a chromosome is duplicated or deleted, it is difficult to identify the specific “target” gene(s) on the chromosome whose gain or loss confers a growth advantage to the tumor cell. Target genes are more easily identified in the case of chromosome translocations, homozygous deletions, and gene amplifications. Translocations generally fuse two genes to create an oncogene (such as *BCR-ABL* in chronic myelogenous leukemia) but, in a small number of cases, can inactivate a tumor suppressor gene by truncating it or separating it from its promoter. Homozygous deletions often involve just one or a few genes, and the target is always a tumor suppressor gene. Amplifications contain an oncogene whose protein product is abnormally active simply because the tumor cell contains 10 to 100 copies of the gene per cell, compared with the two copies present in normal cells.

Most solid tumors have dozens of translocations; however, as with point mutations, the majority of translocations appear to be passengers rather than drivers. The breakpoints of the translocations are often in “gene deserts” devoid of known genes, and many of the translocations and homozygous deletions are adjacent to fragile sites that are prone to breakage. Cancer cells can, perhaps, survive such chromosome breaks more easily than normal cells because they contain mutations that incapacitate genes like *TP53*, which would normally respond to DNA damage by triggering cell death. Studies to date indicate that there are roughly 10 times fewer genes affected by chromosomal changes than by point mutations. Figure 3 shows the types and distribution of genetic alterations that affect protein-coding genes in five representative tumor types. Protein-coding genes account for only ~1.5% of the total genome, and the number of alterations in noncoding regions is proportionately higher than the number affecting coding regions. The vast majority of the alterations in noncoding regions are presumably passengers. These noncoding

mutations, as well as the numerous epigenetic changes found in cancers, will be discussed later.

Drivers Versus Passenger Mutations

Though it is easy to define a “driver gene mutation” in physiologic terms (as one conferring a selective growth advantage), it is more difficult to identify which somatic mutations are drivers and which are passengers. Moreover, it is important to point out that there is a fundamental difference between a driver gene and a driver gene mutation. A driver gene is one that contains driver gene mutations. But driver genes may also contain passenger gene mutations. For example, *APC* is a large driver gene, but only

those mutations that truncate the encoded protein within its N-terminal 1600 amino acids are driver gene mutations. Missense mutations throughout the gene, as well as protein-truncating mutations in the C-terminal 1200 amino acids, are passenger gene mutations.

Numerous statistical methods to identify driver genes have been described. Some are based on the frequency of mutations in an individual gene compared with the mutation frequency of other genes in the same or related tumors after correction for sequence context and gene size (22, 23). Other methods are based on the predicted effects of mutation on the encoded protein, as inferred from biophysical studies (24–26). All of these

methods are useful for prioritizing genes that are most likely to promote a selective growth advantage when mutated. When the number of mutations in a gene is very high, as with *TP53* or *KRAS*, any reasonable statistic will indicate that the gene is extremely likely to be a driver gene. These highly mutated genes have been termed “mountains” (1). Unfortunately, however, genes with more than one, but still relatively few mutations (so called “hills”) numerically dominate cancer genome landscapes (1). In these cases, methods based on mutation frequency and context alone cannot reliably indicate which genes are drivers, because the background rates of mutation vary so much among different patients and regions of the genome. Recent studies of normal cells have indicated that the rate of mutation varies by more than 100-fold within the genome (27). In tumor cells, this variation can be higher and may affect whole




Fig. 3. Total alterations affecting protein-coding genes in selected tumors. Average number and types of genomic alterations per tumor, including single-base substitutions (SBS), small insertions and deletions (indels), amplifications, and homozygous deletions, as determined by genome-wide sequencing studies. For colorectal, breast, and pancreatic ductal cancer, and medulloblastomas, translocations are also included. The published data on which this figure is based are provided in table S1D.

Box 1. Glossary

Adenoma: A benign tumor composed of epithelial cells.

Alternative lengthening of telomeres (ALT): A process of maintaining telomeres independent of telomerase, the enzyme normally responsible for telomere replication.

Amplification: A genetic alteration producing a large number of copies of a small segment (less than a few megabases) of the genome.

Angiogenesis: the process of forming vascular conduits, including veins, arteries, and lymphatics.

Benign tumor: An abnormal proliferation of cells driven by at least one mutation in an oncogene or tumor suppressor gene. These cells are not invasive (i.e., they cannot penetrate the basement membrane lining them), which distinguishes them from malignant cells.

Carcinoma: A type of malignant tumor composed of epithelial cells.

Clonal mutation: A mutation that exists in the vast majority of the neoplastic cells within a tumor.

Driver gene mutation (driver): A mutation that directly or indirectly confers a selective growth advantage to the cell in which it occurs.

Driver gene: A gene that contains driver gene mutations (Mut-Driver gene) or is expressed aberrantly in a fashion that confers a selective growth advantage (Epi-Driver gene).

Epi-driver gene: A gene that is expressed aberrantly in cancers in a fashion that confers a selective growth advantage.

Epigenetic: Changes in gene expression or cellular phenotype caused by mechanisms other than changes in the DNA sequence.

Exome: The collection of exons in the human genome. Exome sequencing generally refers to the collection of exons that encode proteins.

Gatekeeper: A gene that, when mutated, initiates tumorigenesis. Examples include *RB*, mutations of which initiate retinoblastomas, and *VHL*, whose mutations initiate renal cell carcinomas.

Germline genome: An individual's genome, as inherited from their parents.

Germline variants: Variations in sequences observed in different individuals. Two randomly chosen individuals differ by ~20,000 genetic variations distributed throughout the exome.

Human leukocyte antigen (HLA): A protein encoded by genes that determine an individual's capacity to respond to specific antigens or reject transplants from other individuals.

Homozygous deletion: Deletion of both copies of a gene segment (the one inherited from the mother, as well as that inherited from the father).

Indel: A mutation due to small insertion or deletion of one or a few nucleotides.

Karyotype: Display of the chromosomes of a cell on a microscopic slide, used to evaluate changes in chromosome number as well as structural alterations of chromosomes.

Kinase: A protein that catalyzes the addition of phosphate groups to other molecules, such as proteins or lipids. These proteins are essential to nearly all signal transduction pathways.

Liquid tumors: Tumors composed of hematopoietic (blood) cells, such as leukemias. Though lymphomas generally form solid masses in lymph nodes, they are often classified as liquid tumors because of their derivation from hematopoietic cells and ability to travel through lymphatics.

Malignant tumor: An abnormal proliferation of cells driven by mutations in oncogenes or tumor suppressor genes that has already invaded their surrounding stroma. It is impossible to distinguish an isolated benign tumor cell from an isolated malignant tumor cell. This distinction can be made only through examination of tissue architecture.

Metastatic tumor: A malignant tumor that has migrated away from its primary site, such as to draining lymph nodes or another organ.

Methylation: Covalent addition of a methyl group to a protein, DNA, or other molecule.

Missense mutation: A single-nucleotide substitution (e.g., C to T) that results in an amino acid substitution (e.g., histidine to arginine).

Mut-driver gene: A gene that contains driver gene mutations.

Nonsense mutation: A single-nucleotide substitution (e.g., C to T) that results in the production of a stop codon.

Nonsynonymous mutation: A mutation that alters the encoded amino acid sequence of a protein. These include missense, nonsense, splice site, translation start, translation stop, and indel mutations.

Oncogene: A gene that, when activated by mutation, increases the selective growth advantage of the cell in which it resides.

Passenger mutation (passenger): A mutation that has no direct or indirect effect on the selective growth advantage of the cell in which it occurred.

Primary tumor: The original tumor at the site where tumor growth was initiated. This can be defined for solid tumors, but not for liquid tumors.

Promoter: A region within or near the gene that helps regulate its expression.

Rearrangement: A mutation that juxtaposes nucleotides that are normally separated, such as those on two different chromosomes.

Selective growth advantage (*s*): The difference between birth and death in a cell population. In normal adult cells in the absence of injury, $s = 0.000000$.

Self-renewing tissues: Tissues whose cells normally repopulate themselves, such as those lining the gastrointestinal or urogenital tracts, as well as blood cells.

Single-base substitution (SBS): A single-nucleotide substitution (e.g., C to T) relative to a reference sequence or, in the case of somatic mutations, relative to the germline genome of the person with a tumor.

Solid tumors: Tumors that form discrete masses, such as carcinomas or sarcomas.

Somatic mutations: Mutations that occur in any non-germ cell of the body after conception, such as those that initiate tumorigenesis.

Splice sites: Small regions of genes that are juxtaposed to the exons and direct exon splicing.

Stem cell: An immortal cell that can repopulate a particular cell type.

Subclonal mutation: A mutation that exists in only a subset of the neoplastic cells within a tumor.

Translocation: A specific type of rearrangement where regions from two nonhomologous chromosomes are joined.

Tumor suppressor gene: A gene that, when inactivated by mutation, increases the selective growth advantage of the cell in which it resides.

Untranslated regions: Regions within the exons at the 5' and 3' ends of the gene that do not encode amino acids.




Fig. 4. Distribution of mutations in two oncogenes (*PIK3CA* and *IDH1*) and two tumor suppressor genes (*RB1* and *VHL*). The distribution of missense mutations (red arrowheads) and truncating mutations (blue arrowheads) in representative oncogenes and tumor suppressor genes are shown. The data were

collected from genome-wide studies annotated in the COSMIC database (release version 61). For *PIK3CA* and *IDH1*, mutations obtained from the COSMIC database were randomized by the Excel RAND function, and the first 50 are shown. For *RB1* and *VHL*, all mutations recorded in COSMIC are plotted. aa, amino acids.

regions of the genome in an apparently random fashion (28). Thus, at best, methods based on mutation frequency can only prioritize genes for further analysis but cannot unambiguously identify driver genes that are mutated at relatively low frequencies.

Further complicating matters, there are two distinct meanings of the term “driver gene” that are used in the cancer literature. The driver-versus-passenger concept was originally used to distinguish mutations that caused a selective growth advantage from those that did not (29). According to this definition, a gene that does not harbor driver gene mutations cannot be a driver gene. But many genes that contain few or no driver gene mutations have been labeled driver genes in the literature. These include genes that are overexpressed, underexpressed, or epigenetically altered in tumors, or those that enhance or inhibit some aspect of tumorigenicity when their expression is experimentally manipulated. Though a subset of these genes may indeed play an important role in the neoplastic process, it is confusing to lump them all together as driver genes.

To reconcile the two connotations of driver genes, we suggest that genes suspected of increasing the selective growth advantage of tumor cells be categorized as either “Mut-driver genes” or “Epi-driver genes.” Mut-driver genes contain a sufficient number or type of driver gene mutations to unambiguously distinguish them from other genes. Epi-driver genes are expressed aber-

rantly in tumors but not frequently mutated; they are altered through changes in DNA methylation or chromatin modification that persist as the tumor cell divides.

A Ratiometric Method to Identify and Classify Mut-Driven Genes

If mutation frequency, corrected for mutation context, gene length, and other parameters, cannot reliably identify modestly mutated driver genes, what can? In our experience, the best way to identify Mut-driver genes is through their pattern of mutation rather than through their mutation frequency. The patterns of mutations in well-studied oncogenes and tumor suppressor genes are highly characteristic and nonrandom. Oncogenes are recurrently mutated at the same amino acid positions, whereas tumor suppressor genes are mutated through protein-truncating alterations throughout their length (Fig. 4 and table S2A).

On the basis of these mutation patterns rather than frequencies, we can determine which of the 18,306 mutated genes containing a total of 404,863 subtle mutations that have been recorded in the Catalogue of Somatic Mutations in Cancer (COSMIC) database (30) are Mut-driver genes and whether they are likely to function as oncogenes or tumor suppressor genes. To be classified as an oncogene, we simply require that >20% of the recorded mutations in the gene are at recurrent positions and are missense (see legend to table S2A). To be classified as a tumor suppressor

gene, we analogously require that >20% of the recorded mutations in the gene are inactivating. This “20/20 rule” is lenient in that all well-documented cancer genes far surpass these criteria (table S2A).

The following examples illustrate the value of the 20/20 rule. When *IDH1* mutations were first identified in brain tumors, their role in tumorigenesis was unknown (2, 31). Initial functional studies suggested that *IDH1* was a tumor suppressor gene and that mutations inactivated this gene (32). However, nearly all of the mutations in *IDH1* were at the identical amino acid, codon 132 (Fig. 4). As assessed by the 20/20 rule, this distribution unambiguously indicated that *IDH1* was an oncogene rather than a tumor suppressor gene, and this conclusion was eventually supported by biochemical experiments (33, 34). Another example is provided by mutations in *NOTCH1*. In this case, some functional studies suggested that *NOTCH1* was an oncogene, whereas others suggested it was a tumor suppressor gene (35, 36). The situation could be clarified through the application of the 20/20 rule to *NOTCH1* mutations in cancers. In “liquid tumors” such as lymphomas and leukemias, the mutations were often recurrent and did not truncate the predicted protein (37). In squamous cell carcinomas, the mutations were not recurrent and were usually inactivating (38–40). Thus, the genetic data clearly indicated that *NOTCH1* functions differently in different tumor types. The idea that the same gene can function

in completely opposite ways in different cell types is important for understanding cell signaling pathways.

How Many Mut-Driver Genes Exist?

Though all 20,000 protein-coding genes have been evaluated in the genome-wide sequencing studies of 3284 tumors, with a total of 294,881 mutations reported, only 125 Mut-driver genes, as defined by the 20/20 rule, have been discovered to date (table S2A). Of these, 71 are tumor suppressor genes and 54 are oncogenes. An important but relatively small fraction (29%) of these genes was discovered to be mutated through unbiased genome-wide sequencing; most of these genes had already been identified by previous, more directed investigations.

How many more Mut-driver genes are yet to be discovered? We believe that a plateau is being reached, because the same Mut-driver genes keep being “rediscovered” in different tumor types. For example, *MLL2* and *MLL3* mutations were originally discovered in medulloblastomas (41) and were subsequently discovered to be mutated in non-Hodgkin lymphomas, prostate cancers, breast cancers, and other tumor types (42–45). Similarly, *ARID1A* mutations were first discovered to be mutated in clear-cell ovarian cancers (46, 47) and were subsequently shown to be mutated in tumors of several other organs, including those of the stomach and liver (48–50). In recent studies of several types of lung cancer (4, 51, 52), nearly all genes found to be mutated at significant

frequencies had already been identified in tumors of other organs. In other words, the number of frequently altered Mut-driver genes (mountains) is nearing saturation. More mountains will undoubtedly be discovered, but these will likely be in uncommon tumor types that have not yet been studied in depth.

The newly discovered Mut-driver genes that have been detected through genome-wide sequencing have often proved illuminating. For example, nearly half of these genes encode proteins that directly regulate chromatin through modification of histones or DNA. Examples include the histones HIST1H3B and H3F3A, as well as the proteins DNMT1 and TET1, which covalently modify DNA, EZH2, SETD2, and KDM6A, which, in turn, methylate or demethylate histones (53–57). These discoveries have profound implications for understanding the mechanistic basis of the epigenetic changes that are rampant in tumors (58). The discovery of genetic alterations in genes encoding mRNA splicing factors, such as *SF3B1* and *U2AF1* (59–61), was similarly stunning, as mutations in these genes would be expected to lead to a plethora of nonspecific cellular stresses rather than to promote specific tumor types. Another example is provided by mutations in the cooperating proteins ATRX and DAXX (62). Tumors with mutations in these genes all have a specific type of telomere elongation process termed “ALT” (for “alternative lengthening of telomeres”) (63). Though the ALT phenotype had been recognized for more than a decade, its genetic basis

was mysterious before the discovery of mutations of these genes and their perfect correlation with the ALT phenotype (64). A final example is provided by *IDH1* and *IDH2*, whose mutations have stimulated the burgeoning field of tumor metabolism (65) and have had fascinating implications for epigenetics (66, 67).

The Mut-driver genes listed in table S2A are affected by subtle mutations: base substitutions, intragenic insertions, or deletions. As noted above, Mut-driver genes can also be altered by less subtle changes, such as translocations, amplifications, and large-scale deletions. As with point mutations, it can be difficult to distinguish Mut-driver genes that are altered by these types of changes from genes that contain only passenger mutations. Genes that are not point-mutated, but are recurrently amplified (e.g., *MYC* family genes) or homozygously deleted (e.g., *MAP2K4*) and that meet other criteria (e.g., being the only gene in the amplicon or homozygously deleted region) are listed in table S2B. This adds 13 Mut-driver genes—10 oncogenes that are amplified and 3 tumor suppressor genes that are homozygously deleted—to the 125 driver genes that are affected by subtle mutations, for a total of 138 driver genes discovered to date (table S2).

Translocations provide similar challenges for driver classification. An important discovery related to this point is chromothripsis (68), a rare cataclysmic event involving one or a small number of chromosomes that results in a large number of chromosomal rearrangements. This complicates any inferences about causality, in the same way that mismatch repair deficiency compromises the interpretation of point mutations. However, for completeness, all fusion genes that have been identified in at least three independent tumors are listed in table S3. Virtually all of these genes were discovered through conventional approaches before the advent of genome-wide DNA sequencing studies, with some notable exceptions such as those described in (6) and (69). The great majority of these translocations are found in liquid tumors (leukemias and lymphomas) (table S3C) or mesenchymal tumors (table S3B) and were initially identified through karyotypic analyses. A relatively small number of recurrent fusions, the most important of which include *ERG* in prostate cancers (70) and *ALK* in lung cancers (71), have been described in more common tumors (table S3A).

Genes exist that predispose to cancer when inherited in mutant form in the germ line, but are not




Fig. 5. Number and distribution of driver gene mutations in five tumor types. The total number of driver gene mutations [in oncogenes and tumor suppressor genes (TSGs)] is shown, as well as the number of oncogene mutations alone. The driver genes are listed in tables S2A and S2B. Translocations are not included in this figure, because few studies report translocations along with the other types of genetic alterations on a per-case basis. In the tumor types shown here, translocations affecting driver genes occur in less than 10% of samples. The published data on which this figure is based are provided in table S1E.

somatically mutated in cancer to a substantial degree. These genes generally do not confer an increase in selective growth advantage when they are abnormal, but they stimulate tumorigenesis in indirect ways (such as by increasing genetic instability, as discussed later in this Review). For completeness, these genes and the hereditary syndromes for which they are responsible are listed in table S4.

Dark Matter

Classic epidemiologic studies have suggested that solid tumors ordinarily require five to eight “hits,” now interpreted as alterations in driver genes, to develop (72). Is this number compatible with the molecular genetic data? In pediatric tumors such as medulloblastomas, the number of driver gene mutations is low (zero to two), as expected from the discussion above (Fig. 5). In common adult tumors—such as pancreatic, colorectal, breast, and brain cancers—the number of mutated driver genes is often three to six, but several tumors have only one or two driver gene mutations (Fig. 5). How can this be explained, given the widely accepted notion that tumor development and progression require multiple, sequential genetic alterations acquired over decades?

First, technical issues explain some of the “missing mutations.” Genome-wide sequencing is far from perfect, at least with the technologies available today. Some regions of the genome are not well represented because their sequences are difficult to amplify, capture, or unambiguously map to the genome (73–76). Second, there is usually a wide distribution in the number of times that a specific nucleotide in a given gene is observed in the sequence data, so some regions will not be well represented by chance factors alone (77). Finally, primary tumors contain not only neoplastic cells, but also stromal cells that dilute the signal from the mutated base, further reducing the probability of finding a mutation (78).

What fraction of mutations are missed by these three technical issues? A recent study of pancreatic cancers is informative in this regard. Biankin *et al.* used immunohistochemical and genetic analyses to select a set of primary tumor samples enriched in neoplastic cells (79). They used massively parallel sequencing to analyze the exomes of these samples, then compared their mutational data with a set of pancreatic cancer cell lines and xenografts in which mutations had previously been identified, using conventional Sanger sequencing, and confirmed to be present in the primary tumors (3, 16). Only 159 (63%) of the expected 251 driver gene mutations were identified in the primary tumors studied by next-generation sequencing alone, indicating a false-negative rate of 37%. Genome-wide studies in which the proportion of neoplastic cells within tu-

mors is not as carefully evaluated as in (79) will have higher false-negative rates. Moreover, these technical problems are exacerbated in whole-genome studies compared with exomic analyses, because the sequence coverage of the former is often lower than that of the latter (generally 30-fold in whole-genome studies versus more than 100-fold in exomic studies).

Conceptual issues also limit the number of detectable drivers. Virtually all studies, either at the whole-genome or whole-exome level, have focused on the coding regions. The reason for

interpret than the somatic mutations in cancers. The first examples of light coming to such dark matter have recently been published: Recurrent mutations in the promoter of the *TERT* gene, encoding the catalytic subunit of telomerase, have been identified and shown to activate its transcription (81, 82).

Mut-driver genes other than those listed in table S2 will undoubtedly be discovered as genome-wide sequencing continues. However, based on the trends noted above, most of the Mut-driver genes will likely be mountains in




Fig. 6. Four types of genetic heterogeneity in tumors, illustrated by a primary tumor in the pancreas and its metastatic lesions in the liver. Mutations introduced during primary tumor cell growth result in clonal heterogeneity. At the top left, a typical tumor is represented by cells with a large fraction of the total mutations (founder cells) from which subclones are derived. The differently colored regions in the subclones represent stages of evolution within a subclone. **(A)** Intratumoral: heterogeneity among the cells of the primary tumor. **(B)** Intermetastatic: heterogeneity among different metastatic lesions in the same patient. In the case illustrated here, each metastasis was derived from a different subclone. **(C)** Intrametastatic: heterogeneity among the cells of each metastasis develops as the metastases grow. **(D)** Interpatient: heterogeneity among the tumors of different patients. The mutations in the founder cells of the tumors of these two patients are almost completely distinct (see text).

this is practical; it is difficult enough to identify driver gene mutations when they qualitatively alter the sequence of the encoded protein. Trying to make sense of intergenic or intronic mutations is much more difficult. Based on analogous studies of the identifiable mutations in patients with monogenic diseases, more than 80% of mutations should be detectable through analysis of the coding regions (80). However, this still leaves some mutations as unidentifiable “dark matter,” even in the germline genomes of heritable cases, which are usually easier to in-

terpret than the somatic mutations in cancers. The first examples of light coming to such dark matter have recently been published: Recurrent mutations in the promoter of the *TERT* gene, encoding the catalytic subunit of telomerase, have been identified and shown to activate its transcription (81, 82).

Mut-driver genes other than those listed in table S2 will undoubtedly be discovered as genome-wide sequencing continues. However, based on the trends noted above, most of the Mut-driver genes will likely be mountains in

mutation, may confer a selective growth advantage (83, 84).

The most obvious source of dark matter is in Epi-driver genes. Human tumors contain large numbers of epigenetic changes affecting DNA or chromatin proteins. For example, a recent study of colorectal cancers showed that more than 10% of the protein-coding genes were differentially methylated when compared with normal colorectal epithelial cells (85). Some of these changes (i.e., those in Epi-driver genes) are likely to provide a selective growth advantage (86, 87). For example, epigenetic silencing of *CDK2NA* and *MLH1* is much more common than mutational inactivation of either of these two well-recognized driver genes (85). However, there is a critical difference between a genetic and an epigenetic change in a gene. Unlike the sequence of a gene in a given individual, methylation is plastic, varying with cell type, developmental stage, and patient age (21). The methylation state of the normal precursor cells that initiate tumorigenesis is unknown; these cells, such as normal stem cells, may represent only a tiny fraction of the cells in a normal organ. This plasticity also means that methylation can change under microenvironmental cues, such as those associated with low nutrient concentrations or abnormal cell contacts. It is therefore difficult to know whether specific epigenetic changes observed in cancer cells reflect, rather than contribute to, the neoplastic state. Criteria for distinguishing epigenetic changes that exert a selective growth advantage from those that do not (passenger epigenetic changes) have not yet been formulated. Given that Epi-driver genes are likely to compose a major component of the dark matter, further research on this topic is essential (58).

Genetic Heterogeneity

The mutations depicted in Fig. 1 are clonal; that is, they are present in the majority of the neoplastic cells in the tumors. But additional, subclonal (i.e., heterogeneous within the tumor) mutations are important for understanding tumor evolution. Four types of genetic heterogeneity are relevant to tumorigenesis (Fig. 6):

1) Intratumoral: heterogeneity among the cells of one tumor. This type of heterogeneity has been recognized for decades. For example, it is rare to see a cytogenetic study of a solid tumor in which all of the tumor cells display the same karyotype (88). The same phenomenon has been noted for individual genes [e.g., (89)] and more recently has been observed throughout the genome (16, 90–96). This kind of heterogeneity must exist: Every time a normal (or tumor) cell divides, it acquires a few mutations, and the number of mutations that distinguish any two cells simply marks the time from their last common ancestor (their founder cell). Cells at the opposite ends of large tumors will be spa-

tially distinct and, in general, will display more differences than neighboring cells (16). This phenomenon is analogous to speciation, wherein organisms on different islands are more likely to diverge from one another than are organisms on the same island.

In studies that have evaluated intratumoral heterogeneity by genome-wide sequencing, the majority of somatic mutations are present in all tumor cells. These mutations form the trunk of the somatic evolutionary tree. What is the importance of the mutations in the branches (i.e., those that are not shared by all tumor cells)? From a medical perspective, these mutations are often meaningless because the primary tumors are surgically removed. How much heterogeneity existed in the various branches before surgery is not important. However, this heterogeneity provides the seeds for intermetastatic heterogeneity, which is of great clinical importance.

2) Intermetastatic: heterogeneity among different metastatic lesions of the same patient. The vast majority of cancer patients die because their tumors were not removed before metastasis to surgically inaccessible sites, such as the liver, brain, lung, or bone. Patients who relapse with a single metastatic lesion can often still be cured by surgery or radiotherapy, but single metastases are the exception rather than the rule. A typical patient on a clinical trial has a dozen or more metastatic lesions large enough to be visualized by imaging, and many more that are smaller. If each of the metastatic lesions in a single patient was founded by a cell with a very different genetic constitution, then chemotherapeutic cures would be nearly impossible to achieve: Eradicating a subset of the metastatic lesions in a patient will not be adequate for long-term survival.

How much heterogeneity is there among different metastatic lesions? In short, a lot. It is not uncommon for one metastatic lesion to have 20 clonal genetic alterations not shared by other metastases in the same patient (16, 97). Because they are clonal, these mutations occurred in the founder cell of the metastasis; that is, the cell that escaped from the primary tumor and multiplied to form the metastasis. The founder cell for each metastasis is present in different, geographically distinct areas of the primary tumors, as expected (16).

This potentially disastrous situation is tempered by the fact that the heterogeneity appears largely confined to passenger gene mutations. In most of the studies documenting heterogeneity in malignancies, the Mut-driver genes are present in the trunks of the trees, though exceptions have been noted (95). These findings are consistent with the idea, discussed above, that the genetic alterations required for metastasis were present (i.e., selected for) before metastasis actually occurred. The data are also

consistent with the observation that in patients responsive to targeted agents, the response is often seen in all metastatic lesions rather than just a small subset (98).

3) Intrametastatic: heterogeneity among the cells of an individual metastasis. Each metastasis is established by a single cell (or small group of cells) with a set of founder mutations. As it grows, the metastasis acquires new mutations with each cell division. Though the founder mutations may make the lesion susceptible to antitumor agents, the new mutations provide the seeds for drug resistance. Unlike primary tumors, the metastatic lesions generally cannot be removed by surgery and must be treated with systemic therapies. Patients with complete responses to targeted therapies invariably relapse. Most of the initial lesions generally recur, and the time frame at which they recur is notably similar. This time course can be explained by the presence of resistance mutations that existed within each metastasis before the onset of the targeted therapy (99–102). Calculations show that any metastatic lesion of a size visible on medical imaging has thousands of cells (among the billions present) that are already resistant to virtually any drug that can be imagined (99, 101, 102). Thus, recurrence is simply a matter of time, entirely predictable on the basis of known mutation frequencies and tumor cell growth rates. This “fait accompli” can be circumvented, in principle, by treatment with multiple agents, as it is unlikely that a single tumor cell will be resistant to multiple drugs that act on different targets.

4) Interpatient: heterogeneity among the tumors of different patients. This type of heterogeneity has been observed by every oncologist; no two cancer patients have identical clinical courses, with or without therapy. Some of these differences could be related to host factors, such as germline variants that determine drug half-life or vascular permeability to drugs or cells, and some could be related to nongenetic factors (103). However, much of this interpatient heterogeneity is probably related to somatic mutations within tumors. Though several dozen somatic mutations may be present in the breast cancers from two patients, only a small number are in the same genes, and in the vast majority of cases, these are the Mut-driver genes (1, 104, 105). Even in these driver genes, the actual mutations are often different. Mutations altering different domains of a protein would certainly not be expected to have identical effects on cellular properties, as experimentally confirmed (106). Though it may seem that different mutations in adjacent codons would have identical effects, detailed studies of large numbers of patients have shown that this need not be the case. For example, a Gly¹²→Asp¹² (G12D) mutation of *KRAS* does not have the same clinical implications as a G13D mutation of the same gene (107). Interpatient heterogeneity has always been one of the major obstacles

to designing uniformly effective treatments for cancer. Efforts to individualize treatments based on knowledge of the genomes of cancer patients are largely based on an appreciation of this heterogeneity.

Signaling Pathways in Tumors

The immense complexity of cancer genomes that could be inferred from the data described above is somewhat misleading. After all, even advanced tumors are not completely out of control, as evidenced by the dramatic responses to agents that target mutant *BRAF* in melanomas (108) or mutant *ALK* in lung cancers (109). Albeit transient, these responses mean that interference with even a single mutant gene product is sufficient to stop cancer in its tracks, at least transiently. How can the genomic complexity of cancer be reconciled with these clinical observations?

Two concepts bear on this point. The first, mentioned above, is that >99.9% of the alterations in tumors (including point mutations, copy-number alterations, translocations, and epigenetic changes distributed throughout the genome, not just in the coding regions) are immaterial to neoplasia. They are simply passenger changes that mark the time that has elapsed between successive clonal expansions. Normal cells also undergo genetic alterations as they divide, both at the nucleotide and chromosomal levels. However, normal cells are programmed to undergo

cell death in response to such alterations, perhaps as a protective mechanism against cancer. In contrast, cancer cells have evolved to tolerate genome complexity by acquiring mutations in genes such as *TP53* (110). Thus, genomic complexity is, in part, the result of cancer, rather than the cause.

To appreciate the second concept, one must take the 30,000-foot view. A jungle might look chaotic at ground level, but the aerial view shows a clear order, with all the animals gathering at the streams at certain points in the day, and all the streams converging at a river. There is order in cancer, too. Mutations in all of the 138 driver genes listed in table S2 do one thing: cause a selective growth advantage, either directly or indirectly. Moreover, there appears to be only a limited number of cellular signaling pathways through which a growth advantage can be incurred (Fig. 7 and table S5).

All of the known driver genes can be classified into one or more of 12 pathways (Fig. 7). The discovery of the molecular components of these pathways is one of the greatest achievements of biomedical research, a tribute to investigators working in fields that encompass biochemistry, cell biology, and development, as well as cancer. These pathways can themselves be further organized into three core cellular processes:

1) Cell fate: Numerous studies have demonstrated the opposing relationship between cell division and differentiation, the arbiters of cell fate. Dividing cells that are responsible for populating normal tissues (stem cells) do not differentiate, and vice versa. Regenerative medicine is based on this distinction, predicated on ways to get differentiated cells to dedifferentiate into stem cells, then forcing the stem cells to differentiate into useful cell types for transplantation back into the patient. Many of the genetic alterations in cancer abrogate the precise balance between differentiation and division, favoring the latter. This causes a selective growth advantage, because differentiating cells eventually die or become quiescent. Pathways that function through this process include APC, HH, and NOTCH, all of which are well known to control cell fate in organisms ranging from worms to mammals (111). Genes encoding chromatin-modifying enzymes can also be included in this category. In normal development, the heritable switch from division to differentiation is not determined by mutation, as it is in cancer, but rather

by epigenetic alterations affecting DNA and chromatin proteins. What better way to subvert this normal mechanism for controlling tissue architecture than to debilitate the epigenetic modifying apparatus itself?

2) Cell survival: Though cancer cells divide abnormally because of cell-autonomous alterations, such as those controlling cell fate, their surrounding stromal cells are perfectly normal and do not keep pace. The most obvious ramification of this asymmetry is the abnormal vasculature of tumors. As opposed to the well-ordered network of arteries, veins, and lymphatics that control nutrient concentrations in normal tissues, the vascular system in cancers is tortuous and lacks uniformity of structure (112, 113). Normal cells are always within 100 μm of a capillary, but this is not true for cancer cells (114). As a result, a cancer cell acquiring a mutation that allows it to proliferate under limiting nutrient concentrations will have a selective growth advantage, thriving in environments in which its sister cells cannot. Mutations of this sort occur, for example, in the *EGFR*, *HER2*, *FGFR2*, *PDGFR*, *TGFBR2*, *MET*, *KIT*, *RAS*, *RAF*, *PIK3CA*, and *PTEN* genes (table S2A). Some of these genes encode receptors for the growth factors themselves, whereas others relay the signal from the growth factor to the interior of the cell, stimulating growth when activated (115, 116). For instance, mutations in *KRAS* or *BRAF* genes confer on cancer cells the ability to grow in glucose concentrations that are lower than those required for the growth of normal cells or of cancer cells that do not have mutations in these genes (117, 118). Progression through the cell cycle (and its antithesis, apoptosis) can be directly controlled by intracellular metabolites, and driver genes that directly regulate the cell cycle or apoptosis, such as *CDKN2A*, *MYC*, and *BCL2*, are often mutated in cancers. Another gene whose mutations enhance cell survival is *VHL*, the product of which stimulates angiogenesis through the secretion of vascular endothelial growth factor. What better way to provision growth factors to a rogue tumor than to lure the unsuspecting vasculature to its hideout?

3) Genome maintenance: As a result of the exotic microenvironments in which they reside, cancer cells are exposed to a variety of toxic substances, such as reactive oxygen species. Even without microenvironmental poisons, cells make mistakes while replicating their DNA or during division (119, 120), and checkpoints exist to either slow down such cells or make them commit suicide (apoptosis) under such circumstances (110, 121, 122). Although it is good for the organism to remove these damaged cells, tumor cells that can survive the damage will, by definition, have a selective growth advantage. Therefore, it is not surprising that genes whose mutations abrogate these checkpoints, such as *TP53* and *ATM*, are mutated in cancers



Fig. 7. Cancer cell signaling pathways and the cellular processes they regulate. All of the driver genes listed in table S2 can be classified into one or more of 12 pathways (middle ring) that confer a selective growth advantage (inner circle; see main text). These pathways can themselves be further organized into three core cellular processes (outer ring). The publications on which this figure is based are provided in table S5.

(123). Defects in these genes can also indirectly confer a selective growth advantage by allowing cells that have a gross chromosomal change favoring growth, such as a translocation or an extra chromosome, to survive and divide. Analogously, genes that control point mutation rates, such as *MLH1* or *MSH2*, are mutated in cancers (table S2A) or in the germ line of patients predisposed to cancers (table S4) because they accelerate the acquisition of mutations that function through processes that regulate cell fate or survival. What better way to promote cancer than by increasing the rate of occurrence of the mutations that drive the process?

Because the protein products of genes regulating cell fate, cell survival, and genome maintenance often interact with one another, the pathways within them overlap; they are not as discrete as might be inferred from the description above. However, grouping genes into pathways makes perfect sense from a genetics standpoint. Given that cancer is a genetic disease, the principles of genetics should apply to its pathogenesis. When performing a conventional mutagenesis screen in bacteria, yeast, fruit flies, or worms, one expects to discover mutations in several different genes that confer similar phenotypes. The products of these genes often interact with one another and define a biochemical or developmental pathway. Therefore, it should not be surprising that several different genes can result in the same selective growth advantage for cancer cells and that the products of these genes interact. The analogy between cancer pathways and biochemical or developmental pathways in other organisms goes even deeper: The vast majority of our knowledge of the function of driver genes has been derived from the study of the pathways through which their homologs work in nonhuman organisms. Though the functions are not identical to those in human cells, they are highly related and have provided the starting point for analogous studies in human cells.

Recognition of these pathways also has important ramifications for our ability to understand interpatient heterogeneity. One lung cancer might have an activating mutation in a receptor for a stimulatory growth factor, making it able to grow in low concentrations of epidermal growth factor (EGF). A second lung cancer might have an activating mutation in *KRAS*, whose protein product normally transmits the signal from the epidermal growth factor receptor (EGFR) to other cell signaling molecules. A third lung cancer might have an inactivating mutation in *NF1*, a regulatory protein that normally inactivates the *KRAS* protein. Finally, a fourth lung cancer might have a mutation in *BRAF*, which transmits the signal from *KRAS* to downstream kinases (Fig. 8). One would predict that mutations in the various components of a single pathway would be mutually exclusive—that is, not occurring in the




Fig. 8. Signal transduction pathways affected by mutations in human cancer. Two representative pathways from Fig. 7 (RAS and PI3K) are illustrated. The signal transducers are color coded: red indicates protein components encoded by the driver genes listed in table S2; yellow balls denote sites of phosphorylation. Examples of therapeutic agents that target some of the signal transducers are shown. RTK, receptor tyrosine kinase; GDP, guanosine diphosphate; MEK, MAPK kinase; ERK, extracellular signal-regulated kinase; NFkB, nuclear factor κ B; mTOR, mammalian target of rapamycin.

same tumor—and this has been experimentally confirmed (124, 125). Apart from being intellectually satisfying, knowledge of these pathways has implications for cancer therapy, as discussed in the next section.

A Perspective on Genome-Based Medicine in Oncology

Opportunities

Though cancer genome sequencing is a relatively new endeavor, it has already had an impact on the

clinical care of cancer patients. The recognition that certain tumors contain activating mutations in driver genes encoding protein kinases has led to the development of small-molecule inhibitor drugs targeting those kinases.

Representative examples of this type of genome-based medicine include the use of EGFR kinase inhibitors to treat cancers with *EGFR* gene mutations (126), the aforementioned anaplastic lymphoma kinase (ALK) inhibitors to treat cancers with *ALK* gene translocations (109), and specific inhibitors of mutant BRAF

to treat cancers with *BRAF* mutations (108). Before instituting treatment with such agents, it is imperative to determine whether the cancer harbors the mutations that the drug targets. Only a small fraction of lung cancer patients have *EGFR* gene mutations or *ALK* gene translocations, and only these patients will respond to the drugs. Treating lung cancer patients without these particular genetic alterations would be detrimental, as such patients would develop the toxic side effects of the drugs while their tumors progressed.

A second type of genome-based medicine focuses on the side effects and metabolism of the therapeutic agents, rather than the genetic alterations they target. At present, the dose of cancer drugs given to patients is based on the patients' size (body weight or surface area). But the therapeutic ratio of cancer drugs (ratio of the concentration that causes side effects to the concentration required to kill tumor cells) is generally low, particularly for conventional (nontargeted) therapeutic agents. Small changes in circulating concentrations of these drugs can make the difference between substantial tumor regression and intolerable side effects. Interrogation of the germline status of the genes encoding drug-metabolizing enzymes could substantially improve the outcomes of treatment by informing drug dosing (127). Optimally, this genome interrogation would be accompanied by pharmacokinetic measurements of drug concentrations in each patient. The additional cost of such analyses would be small compared with the exorbitant costs of new cancer therapies—for recently approved drugs, the cost is estimated to be \$200,000 to \$300,000 per quality life year produced (128).

Challenges

One challenge of genome-based medicine in oncology is already apparent from the opportunities described above: All of the clinically approved drugs that target the products of genetically altered genes are directed against kinases. One reason for this is that kinases are relatively easy to target with small molecules and have been extensively studied at the biochemical, structural, and physiologic levels (129). But another reason has far deeper ramifications. The vast majority of drugs on the market today, for cancer or other diseases, inhibit the actions of their protein targets. This inhibition occurs because the drugs interfere with the protein's enzymatic activity (such as the phosphorylation catalyzed by kinases) or with the binding of the protein to a small ligand (such as with G protein-coupled receptors). Only 31 of the oncogenes listed in tables S2 and S3 have enzymatic activities that are targetable in this manner. Many others participate in protein complexes, involving large interfaces and numerous weak interactions. Inhibiting the function of such proteins

with small drugs is notoriously difficult because small compounds can only inhibit one of these interactions (130, 131).

Though one can at least imagine the development of drugs that inhibit nonenzymatic protein functions, the second challenge evident from table S2 poses even greater difficulties: A large fraction of the Mut-driver genes encode tumor suppressors. Drugs generally interfere with protein function; they cannot, in general, replace the function of defective genes such as those resulting from mutations in tumor suppressor genes. Unfortunately, tumor suppressor gene-inactivating mutations predominate over oncogene-activating mutations in the most common solid tumors: Few individual tumors contain more than one oncogene mutation (Fig. 5).

The relatively small number of oncogene mutations in tumors is important in light of the intrametastatic heterogeneity described earlier. To circumvent the inevitable development of resistance to targeted therapies, it will likely be necessary to treat patients with two or more drugs. The probability that a single cancer cell within a large metastatic lesion will be resistant to two agents that target two independent pathways is exponentially less than the probability that the cell will be resistant to a single agent. However, if the cancer cell does not contain more than one targetable genetic alteration (i.e., an oncogene mutation), then this combination strategy is not feasible.

Given the paucity of oncogene alterations in common solid tumors and these principles, can

targeted therapeutic approaches ever be expected to induce long-term remissions, even cures, rather than the short-term remissions now being achieved? The saviors are pathways; every tumor suppressor gene inactivation is expected to result in the activation of some growth-promoting signal downstream of the pathway. An example is provided by *PTEN* mutations: Inactivation of the tumor suppressor gene *PTEN* results in activation of the AKT kinase (Fig. 8). Similarly, inactivation of the tumor suppressor gene *CDKN2A* results in activation of kinases, such as cyclin-dependent kinase 4, that promote cell cycle traverse (132). Furthermore, inactivation of tumor suppressor gene *APC* results in constitutive activity of oncogenes such as *CTNNB1* and *CMYC* (133–135).

We believe that greater knowledge of these pathways and the ways in which they function is the most pressing need in basic cancer research. Successful research on this topic should allow the development of agents that target, albeit indirectly, defective tumor suppressor genes. Indeed, there are already examples of such indirect targeting. Inactivating mutations of the tumor suppressor genes *BRCA1* or *BRCA2* lead to activation of downstream pathways required to repair DNA damage in the absence of BRCA function. Thus, cancer cells with defects in *BRCA1* or *BRCA2* are more susceptible to DNA damaging agents or to drugs that inhibit enzymes that facilitate the repair of DNA damage such as PARP [poly(adenosine diphosphate-ribose) polymerase] (136). PARP inhibitors have shown

Box 2. Highlights

1. Most human cancers are caused by two to eight sequential alterations that develop over the course of 20 to 30 years.
2. Each of these alterations directly or indirectly increases the ratio of cell birth to cell death; that is, each alteration causes a selective growth advantage to the cell in which it resides.
3. The evidence to date suggests that there are ~140 genes whose intragenic mutations contribute to cancer (so-called Mut-driver genes). There are probably other genes (Epi-driver genes) that are altered by epigenetic mechanisms and cause a selective growth advantage, but the definitive identification of these genes has been challenging.
4. The known driver genes function through a dozen signaling pathways that regulate three core cellular processes: cell fate determination, cell survival, and genome maintenance.
5. Every individual tumor, even of the same histopathologic subtype as another tumor, is distinct with respect to its genetic alterations, but the pathways affected in different tumors are similar.
6. Genetic heterogeneity among the cells of an individual tumor always exists and can impact the response to therapeutics.
7. In the future, the most appropriate management plan for a patient with cancer will be informed by an assessment of the components of the patient's germline genome and the genome of his or her tumor.
8. The information from cancer genome studies can also be exploited to improve methods for prevention and early detection of cancer, which will be essential to reduce cancer morbidity and mortality.

encouraging results in clinical trials when used in patients whose tumors have inactivating mutations of BRCA genes (137).

Further progress in this area will require more detailed information about the signaling pathways through which cancer genes function in human cancer cells, as well as in model organisms. One of the lessons of molecular biology over the past two decades is that pathway functions are different, depending on the organism, cell type, and precise genetic alterations in that cell (138). A pertinent example of this principle is provided by results of treatment with drugs inhibiting mutant BRAF kinase activity. In the majority of patients with melanomas harboring (V600E; V, Val; E, Glu) mutations in the BRAF gene, these drugs induce dramatic (though transient) remissions (108). But the same drugs have no therapeutic effect in colorectal cancer patients harboring the identical BRAF mutations (139). This observation has been attributed to the expression of EGFR, which occurs in some colorectal cancers but not in melanoma and is thought to circumvent the growth-inhibitory effects of the BRAF inhibitors. With this example in mind, no one should be surprised that a new drug that works well in an engineered tumor in mice fails in human trials; the organism is different, the cell type is usually different, and the precise genetic constitutions are always different. The converse of this statement—that a drug that fails in animal trials will not necessarily fail in human trials—has important practical consequences. In our view, if the biochemical and conceptual bases for a drug's actions are solid and the drug is shown to be safe in animals, then a human trial may be warranted, even if it does not shrink tumors in mice.

Genome-Based Medicines of the Future

Cancer genomes can also be exploited for the development of more effective immunotherapies. As noted above, typical solid tumors contain 30 to 70 mutations that alter the amino acid sequences of the proteins encoded by the affected genes. Each of these alterations is foreign to the immune system, as none have been encountered during embryonic or postnatal life. Therefore, these alterations, in principle, provide a “holy grail” for tumor immunology: truly tumor-specific antigens. These antigens could be incorporated into any of the numerous platforms that already exist for the immunotherapy of cancer. These include administration of vaccines containing the mutant peptide, viruses encoding the mutant peptides on their surfaces, dendritic cells presenting the mutated peptide, and antibodies or T cells with reactivity directed against the mutant peptides (140).

To realize these sorts of therapeutics, several conditions must be met. First, the mutant protein must be expressed. As cancer cells generally express about half of the proteins that are encoded

by the human genome (141), this condition is not limiting. Second, as most proteins affected by mutations are intracellular, these mutations will not be visible to the immune system unless the mutant residue is presented in the context of a human leukocyte antigen (HLA) protein. Based on in silico analyses of binding affinities, it has been estimated that a typical breast or colorectal cancer contains 7 to 10 mutant proteins that can bind to an individual patient's HLA type (142). These theoretical predictions have recently gained experimental support. Studies of mouse tumors have identified mutant genes and shown that the corresponding peptides can induce antitumor immunity when administered as vaccines (143). Moreover, clinical trials of brain cancer patients immunized against a mutant peptide have yielded encouraging results (144).

As with all cancer therapies that are attractive in concept, obstacles abound in practice. If a tumor expresses a mutant protein that is recognizable as foreign, why has the host immune system not eradicated that tumor already? Indeed, immunoediting in cancers has been shown to exist, resulting in the down-regulation or absence of mutant epitopes that should have, and perhaps did, elicit an immune response during tumor development (145, 146). Additionally, tumors can lose immunogenicity through a variety of genetic alterations, thereby precluding the presentation of epitopes that would otherwise be recognized as foreign (147). Though these theoretical limitations are disheartening, recent studies on immune regulation in humans portend cautious optimism (148, 149).

Other Ways to Reduce Morbidity and Mortality Through Knowledge of Cancer Genomics

When we think about eradicating cancer, we generally think about curing advanced cases—those that cannot be cured by surgery alone because they have already metastasized. This is a curious way of thinking about this disease. When we think of cardiovascular or infectious diseases, we first consider ways to prevent them rather than drugs to cure their most advanced forms. Today, we are in no better position to cure polio or massive myocardial infarctions than we were a thousand years ago. But we can prevent these diseases entirely (vaccines), reduce incidence (dietary changes, statins), or mitigate severity (stents, thrombolytic agents) and thereby make a major impact on morbidity and mortality.

This focus on curing advanced cancers might have been reasonable 50 years ago, when the molecular pathogenesis of cancers was mysterious and when chemotherapeutic agents against advanced cancers were showing promise. But this mindset is no longer acceptable. We now know precisely what causes cancer: a sequential series of alterations in well-defined genes that

alter the function of a limited number of pathways. Moreover, we know that this process takes decades to develop and that the incurable stage, metastasis, occurs only a few years before death. In other words, of the one million people that will die from cancer this year, the vast majority will die only because their cancers were not detected in the first 90% of the cancers' lifetimes, when they were amenable to the surgeons' scalpel.

This new knowledge of cancer (Box 2) has reinvigorated the search for cures for advanced cancers, but has not yet permeated other fields of applied cancer research. A common and limited set of driver genes and pathways is responsible for most common forms of cancer (table S2); these genes and pathways offer distinct potential for early diagnosis. The genes themselves, the proteins encoded by these genes, and the end products of their pathways are, in principle, detectable in many ways, including analyses of relevant body fluids, such as urine for genitourinary cancers, sputum for lung cancers, and stool for gastrointestinal cancers (150). Equally exciting are the possibilities afforded by molecular imaging, which not only indicate the presence of a cancer but also reveal its precise location and extent. Additionally, research into the relationship between particular environmental influences (diet and lifestyle) and the genetic alterations in cancer is sparse, despite its potential for preventative measures.

The reasons that society invests so much more in research on cures for advanced cancers than on prevention or early detection are complex. Economic issues play a part: New drugs are far more lucrative for industry than new tests, and large individual costs for treating patients with advanced disease have become acceptable, even in developing countries (151). From a technical standpoint, the development of new and improved methods for early detection and prevention will not be easy, but there is no reason to assume that it will be more difficult than the development of new therapies aimed at treating widely metastatic disease.

Our point is not that strenuous efforts to develop new therapies for advanced cancer patients should be abandoned. These will always be required, no matter our arsenal of early detection or preventative measures. Instead, we are suggesting that “plan A” should be prevention and early detection, and “plan B” (therapy for advanced cancers) should be necessary only when plan A fails. To make plan A viable, government and philanthropic organizations must dedicate a much greater fraction of their resources to this cause, with long-term considerations in mind. We believe that cancer deaths can be reduced by more than 75% in the coming decades (152), but that this reduction will only come about if greater efforts are made toward early detection and prevention.

References and Notes

1. L. D. Wood *et al.*, *Science* **318**, 1108 (2007).
2. D. W. Parsons *et al.*, *Science* **321**, 1807 (2008).
3. S. Jones *et al.*, *Science* **321**, 1801 (2008).
4. R. Govindan *et al.*, *Cell* **150**, 1121 (2012).
5. R. Gryfe, S. Gallinger, *Surgery* **130**, 17 (2001).
6. The Cancer Genome Atlas Network, *Nature* **487**, 330 (2012).
7. C. Palles *et al.*, *Nat. Genet.* **45**, 136 (2013).
8. P. C. Nowell, *Science* **194**, 23 (1976).
9. E. R. Fearon, B. Vogelstein, *Cell* **61**, 759 (1990).
10. K. W. Kinzler, B. Vogelstein, *Nature* **386**, 761 (1997).
11. S. Jones *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **105**, 4283 (2008).
12. I. Bozic *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **107**, 18545 (2010).
13. C. Tomasetti, B. Vogelstein, G. Parmigiani, *Proc. Natl. Acad. Sci. U.S.A.* **110**, 1999 (2013).
14. E. Laurenti, J. E. Dick, *Ann. N.Y. Acad. Sci.* **1266**, 68 (2012).
15. J. S. Welch *et al.*, *Cell* **150**, 264 (2012).
16. S. Yachida *et al.*, *Nature* **467**, 1114 (2010).
17. R. S. Kerbel, *Adv. Cancer Res.* **55**, 87 (1990).
18. R. Bernards, R. A. Weinberg, *Nature* **418**, 823 (2002).
19. M. Yu, S. Stott, M. Toner, S. Maheswaran, D. A. Haber, *J. Cell Biol.* **192**, 373 (2011).
20. J. Komori, L. Boone, A. DeWard, T. Hoppe, E. Lagasse, *Nat. Biotechnol.* **30**, 976 (2012).
21. M. Pelizzola, J. R. Ecker, *FEBS Lett.* **585**, 1994 (2011).
22. G. Parmigiani *et al.*, *Genomics* **93**, 17 (2009).
23. M. Meyerson, S. Gabriel, G. Getz, *Nat. Rev. Genet.* **11**, 685 (2010).
24. H. Carter *et al.*, *Cancer Res.* **69**, 6660 (2009).
25. A. Youn, R. Simon, *Bioinformatics* **27**, 175 (2011).
26. J. S. Kaminker, Y. Zhang, C. Watanabe, Z. Zhang, *Nucleic Acids Res.* **35**, W595 (2007).
27. J. J. Michaelson *et al.*, *Cell* **151**, 1431 (2012).
28. S. Nik-Zainal *et al.*, *Cell* **149**, 979 (2012).
29. S. Thiagalingam *et al.*, *Nat. Genet.* **13**, 343 (1996).
30. S. A. Forbes *et al.*, *Nucleic Acids Res.* **39**, D945 (2011).
31. H. Yan *et al.*, *N. Engl. J. Med.* **360**, 765 (2009).
32. S. Zhao *et al.*, *Science* **324**, 261 (2009).
33. P. S. Ward *et al.*, *Cancer Cell* **17**, 225 (2010).
34. L. Dang *et al.*, *Nature* **465**, 966 (2010).
35. W. S. Pear, J. C. Aster, *Curr. Opin. Hematol.* **11**, 426 (2004).
36. M. Nicolas *et al.*, *Nat. Genet.* **33**, 416 (2003).
37. A. P. Weng *et al.*, *Science* **306**, 269 (2004).
38. N. Agrawal *et al.*, *Science* **333**, 1154 (2011).
39. N. Stransky *et al.*, *Science* **333**, 1157 (2011).
40. N. Agrawal *et al.*, *Cancer Discovery* **2**, 899 (2012).
41. D. W. Parsons *et al.*, *Science* **331**, 435 (2011).
42. L. Pasqualucci *et al.*, *Nat. Genet.* **43**, 830 (2011).
43. R. D. Morin *et al.*, *Nature* **476**, 298 (2011).
44. C. S. Grasso *et al.*, *Nature* **487**, 239 (2012).
45. M. J. Ellis *et al.*, *Nature* **486**, 353 (2012).
46. K. C. Wiegand *et al.*, *N. Engl. J. Med.* **363**, 1532 (2010).
47. S. Jones *et al.*, *Science* **330**, 228 (2010).
48. S. Jones *et al.*, *Hum. Mutat.* **33**, 100 (2012).
49. K. Wang *et al.*, *Nat. Genet.* **43**, 1219 (2011).
50. J. Huang *et al.*, *Nat. Genet.* **44**, 1117 (2012).
51. M. Imielinski *et al.*, *Cell* **150**, 1107 (2012).
52. C. M. Rudin *et al.*, *Nat. Genet.* **44**, 1111 (2012).
53. F. Delhommeau *et al.*, *N. Engl. J. Med.* **360**, 2289 (2009).
54. J. Schwartzentruber *et al.*, *Nature* **482**, 226 (2012).
55. G. Wu *et al.*, *Nat. Genet.* **44**, 251 (2012).
56. T. J. Ley *et al.*, *N. Engl. J. Med.* **363**, 2424 (2010).
57. G. L. Dalglish *et al.*, *Nature* **463**, 360 (2010).
58. M. L. Suvà, N. Riggi, B. E. Bernstein, *Science* **339**, 1567 (2013).
59. E. Papaemmanuil *et al.*, *N. Engl. J. Med.* **365**, 1384 (2011).
60. T. A. Graubert *et al.*, *Nat. Genet.* **44**, 53 (2012).
61. K. Yoshida *et al.*, *Nature* **478**, 64 (2011).
62. Y. Jiao *et al.*, *Science* **331**, 1199 (2011).
63. A. J. Cesare, R. R. Reddel, *Nat. Rev. Genet.* **11**, 319 (2010).
64. C. M. Heaphy *et al.*, *Science* **333**, 425 (2011).
65. M. G. Vander Heiden, L. C. Cantley, C. B. Thompson, *Science* **324**, 1029 (2009).
66. C. Lu *et al.*, *Nature* **483**, 474 (2012).
67. S. Turcan *et al.*, *Nature* **483**, 479 (2012).
68. P. J. Stephens *et al.*, *Cell* **144**, 27 (2011).
69. A. J. Bass *et al.*, *Nat. Genet.* **43**, 964 (2011).
70. S. A. Tomlins *et al.*, *Science* **310**, 644 (2005).
71. M. Soda *et al.*, *Nature* **448**, 561 (2007).
72. P. Armitage, R. Doll, *Br. J. Cancer* **8**, 1 (1954).
73. R. Böttcher *et al.*, *Nucleic Acids Res.* **40**, e125 (2012).
74. M. Forster *et al.*, *Nucleic Acids Res.* **41**, e16 (2013).
75. J. Ding *et al.*, *Bioinformatics* **28**, 167 (2012).
76. M. A. Beal, T. C. Glenn, C. M. Somers, *Mutat. Res.* **750**, 96 (2012).
77. F. Mertes *et al.*, *Brief. Funct. Genomics* **10**, 374 (2011).
78. M. Gundry, J. Vijg, *Mutat. Res.* **729**, 1 (2012).
79. A. V. Biankin *et al.*, *Nature* **491**, 399 (2012).
80. H. Yan, K. W. Kinzler, B. Vogelstein, *Science* **289**, 1890 (2000).
81. F. W. Huang *et al.*, *Science* **339**, 957 (2013).
82. S. Horn *et al.*, *Science* **339**, 959 (2013).
83. W. Xue *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **109**, 8212 (2012).
84. N. L. Solimini *et al.*, *Science* **337**, 104 (2012).
85. A. D. Beggs *et al.*, *J. Pathol.* **10.1002/path.4132** (2012).
86. A. P. Feinberg, B. Tycko, *Nat. Rev. Cancer* **4**, 143 (2004).
87. P. A. Jones, S. B. Baylin, *Cell* **128**, 683 (2007).
88. M. Höglund, D. Gisselsson, T. Säll, F. Mitelman, *Cancer Genet. Cytogenet.* **135**, 103 (2002).
89. D. Shibata, J. Schaeffer, Z. H. Li, G. Capella, M. Perucho, *J. Natl. Cancer Inst.* **85**, 1058 (1993).
90. A. Sottoriva, I. Spiteri, D. Shibata, C. Curtis, S. Tavaré, *Cancer Res.* **73**, 41 (2013).
91. S. P. Shah *et al.*, *Nature* **461**, 809 (2009).
92. K. Anderson *et al.*, *Nature* **469**, 356 (2011).
93. N. Navin *et al.*, *Nature* **472**, 90 (2011).
94. S. Nik-Zainal *et al.*, *Cell* **149**, 994 (2012).
95. M. Gerlinger *et al.*, *N. Engl. J. Med.* **366**, 883 (2012).
96. X. Xu *et al.*, *Cell* **148**, 886 (2012).
97. P. J. Campbell *et al.*, *Nature* **467**, 1109 (2010).
98. N. Wagle *et al.*, *J. Clin. Oncol.* **29**, 3085 (2011).
99. N. L. Komarova, D. Wodarz, *Proc. Natl. Acad. Sci. U.S.A.* **102**, 9714 (2005).
100. A. B. Turke *et al.*, *Cancer Cell* **17**, 77 (2010).
101. R. Durrett, S. Moseley, *Theor. Popul. Biol.* **77**, 42 (2010).
102. L. A. Diaz Jr. *et al.*, *Nature* **486**, 537 (2012).
103. A. Kreso *et al.*, *Science* **339**, 543 (2013).
104. P. J. Stephens *et al.*, *Nature* **486**, 400 (2012).
105. The Cancer Genome Atlas Network, *Nature* **490**, 61 (2012).
106. H. Pang *et al.*, *Cancer Res.* **69**, 8868 (2009).
107. J. Chen, Y. Ye, H. Sun, G. Shi, *Cancer Chemother. Pharmacol.* **71**, 265 (2013).
108. P. B. Chapman *et al.*, *N. Engl. J. Med.* **364**, 2507 (2011).
109. E. L. Kwak *et al.*, *N. Engl. J. Med.* **363**, 1693 (2010).
110. M. Ljungman, D. P. Lane, *Nat. Rev. Cancer* **4**, 727 (2004).
111. N. Perrimon, C. Pitsouli, B. Z. Shilo, *Cold Spring Harb. Perspect. Biol.* **4**, a005975 (2012).
112. R. S. Kerbel, *N. Engl. J. Med.* **358**, 2039 (2008).
113. A. S. Chung, N. Ferrara, *Annu. Rev. Cell Dev. Biol.* **27**, 563 (2011).
114. J. W. Baish *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **108**, 1799 (2011).
115. N. E. Hynes, H. A. Lane, *Nat. Rev. Cancer* **5**, 341 (2005).
116. N. Turner, R. Grose, *Nat. Rev. Cancer* **10**, 116 (2010).
117. J. Yun *et al.*, *Science* **325**, 1555 (2009).
118. H. Ying *et al.*, *Cell* **149**, 656 (2012).
119. D. J. Araten *et al.*, *Cancer Res.* **65**, 8111 (2005).
120. T. A. Kunkel, *Cold Spring Harb. Symp. Quant. Biol.* **74**, 91 (2009).
121. B.-B. S. Zhou, S. J. Elledge, *Nature* **408**, 433 (2000).
122. R. H. Medema, L. Macürek, *Oncogene* **31**, 2601 (2012).
123. F. A. Derheimer, M. B. Kastan, *FEBS Lett.* **584**, 3675 (2010).
124. G. Ciriello, E. Cerami, C. Sander, N. Schultz, *Genome Res.* **22**, 398 (2012).
125. C. H. Yeang, F. McCormick, A. Levine, *FASEB J.* **22**, 2605 (2008).
126. S. V. Sharma, D. W. Bell, J. Settleman, D. A. Haber, *Nat. Rev. Cancer* **7**, 169 (2007).
127. H. L. McLeod, *Science* **339**, 1563 (2013).
128. D. W. Brock, *Oncologist* **15** (suppl. 1), 36 (2010).
129. M. A. Lemmon, J. Schlessinger, *Cell* **141**, 1117 (2010).
130. M. R. Arkin, J. A. Wells, *Nat. Rev. Drug Discov.* **3**, 301 (2004).
131. R. J. Binstock, *Curr. Pharm. Des.* **18**, 1240 (2012).
132. A. Besson, S. F. Dowdy, J. M. Roberts, *Dev. Cell* **14**, 159 (2008).
133. P. J. Morin *et al.*, *Science* **275**, 1787 (1997).
134. T. C. He *et al.*, *Science* **281**, 1509 (1998).
135. M. van de Wetering *et al.*, *Cell* **111**, 241 (2002).
136. H. Farmer *et al.*, *Nature* **434**, 917 (2005).
137. S. Irshad, A. Ashworth, A. Tutt, *Expert Rev. Anticancer Ther.* **11**, 1243 (2011).
138. D. A. Gruneberg *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* **105**, 16472 (2008).
139. M. Mao *et al.*, *Clin. Cancer Res.* **19**, 657 (2013).
140. J. M. Kirkwood *et al.*, *CA Cancer J. Clin.* **62**, 309 (2012).
141. T. Barrett *et al.*, *Nucleic Acids Res.* **39**, D1005 (2011).
142. N. H. Segal *et al.*, *Cancer Res.* **68**, 889 (2008).
143. J. C. Castle *et al.*, *Cancer Res.* **72**, 1081 (2012).
144. J. H. Sampson *et al.*, *J. Clin. Oncol.* **28**, 4722 (2010).
145. H. Matsushita *et al.*, *Nature* **482**, 400 (2012).
146. M. DuPage, C. Mazumdar, L. M. Schmidt, A. F. Cheung, T. Jacks, *Nature* **482**, 405 (2012).
147. M. Challal-Malladi *et al.*, *Cancer Cell* **20**, 728 (2011).
148. F. S. Hodi *et al.*, *N. Engl. J. Med.* **363**, 711 (2010).
149. S. L. Topalian *et al.*, *N. Engl. J. Med.* **366**, 2443 (2012).
150. B. K. Dunn, K. Jegalian, P. Greenwald, *Recent Results Cancer Res.* **188**, 21 (2011).
151. L. C. Lopes, S. Barberato-Filho, A. C. Costa, C. G. S. Osorio-de-Castro, *Rev. Saude Publica* **44**, 620 (2010).
152. G. A. Colditz, K. Y. Wolin, S. Gehlert, *Sci. Transl. Med.* **4**, 127r46 (2012).

Acknowledgments: We thank M. Nowak and I. Bozic for critical reading of the manuscript, S. Gabelli for assisting with the production of Fig. 8, and A. Dixon, V. Ferranta, and E. Cook for artwork. This work was supported by The Virginia and D.K. Ludwig Fund for Cancer Research; The Lustgarten Foundation for Pancreatic Cancer Research; and NIH grants CA 43460, CA 47345, CA 62924, and CA 121113. All authors are Founding Scientific Advisors of Personal Genome Diagnostics (PGDx), a company focused on the identification of genetic alterations in human cancer for diagnostic and therapeutic purposes. All authors are also members of the Scientific Advisory Board of Inostics, a company that is developing technologies for the molecular diagnosis of cancer. All authors own stock in PGDx and Inostics. The terms of these arrangements are being managed by Johns Hopkins University, in accordance with their conflict-of-interest policies.

Supplementary Materials
 www.sciencemag.org/cgi/content/full/339/6127/1546/DC1
 Tables S1 to S5
 10.1126/science.1235122