



# The Role of Magnetic Resonance Imaging (MRI) in the Differentiation of Necrotic Neoplastic Lesions and Abscesses Using Multiparametric Techniques.

Mr. Ibrahim Ado<sup>1</sup>, Dr. Prasad Hedge<sup>2</sup>, Dr. Rajesh Yadav<sup>3</sup>,

<sup>1</sup> Student, <sup>2</sup> Professor and Head Department of Radio diagnosis<sup>2</sup>, Assistance Professor and Department of Radio diagnosis<sup>3</sup>

<sup>1</sup> Department of radio diagnostic,

<sup>1</sup> Nims University, Jaipur Rajasthan 303121

## Abstract

Magnetic Resonance Imaging (MRI) plays a pivotal role in distinguishing between necrotic neoplastic lesions and abscesses, two conditions that can present similarly on conventional imaging sequences. Misdiagnosis of these conditions could lead to inappropriate management, as the treatment strategies for each differ significantly. This study explores the utility of multiparametric MRI, combining conventional imaging sequences with advanced functional imaging techniques, to improve the accuracy of this differentiation. We analyzed the MRI features of necrotic tumors and abscesses in 50 patients using Diffusion-Weighted Imaging (DWI), Dynamic Contrast-Enhanced MRI (DCE-MRI), and Magnetic Resonance Spectroscopy (MRS). Our results demonstrated that the combination of these advanced imaging techniques significantly improved diagnostic accuracy. Multiparametric MRI achieved 92% sensitivity and 89% specificity for differentiating necrotic neoplastic lesions from abscesses, outperforming conventional MRI (68% sensitivity, 70% specificity). The inclusion of DWI, DCE-MRI, and MRS enhanced lesion characterization by assessing diffusion, perfusion, and metabolic features, thereby providing more reliable differentiation than structural imaging alone. These findings suggest that multiparametric MRI can be a powerful tool for improving diagnostic precision in distinguishing these clinically important lesions, thereby aiding in appropriate management and treatment decisions.

## Keywords:

MRI, Necrotic Neoplastic Lesions, Abscesses, Diffusion-Weighted Imaging, Dynamic Contrast-Enhanced MRI, Magnetic Resonance Spectroscopy, Tumor Differentiation, Abscess Diagnosis.

## **Introduction**

The differentiation of necrotic neoplastic lesions from abscesses is a complex challenge in clinical radiology<sup>1</sup>. Both conditions can present with similar imaging features, including fluid-filled regions with peripheral enhancement. However, the management of these two conditions is vastly different: abscesses typically require drainage and antibiotic therapy, while necrotic neoplastic lesions may require surgical resection, chemotherapy, or radiotherapy. Correct diagnosis, therefore, is crucial<sup>4-6</sup>.

Conventional MRI sequences (T1- and T2-weighted imaging) can provide some clues regarding lesion morphology and enhancement patterns but often lack the specificity needed for definitive diagnosis. Therefore, multiparametric MRI—which combines various advanced MRI techniques such as Diffusion-Weighted Imaging (DWI), Dynamic Contrast-Enhanced MRI (DCE-MRI), and Magnetic Resonance Spectroscopy (MRS)—has emerged as an essential tool for more accurate diagnosis. These techniques provide functional and metabolic data that supplement the anatomical information from conventional MRI<sup>9-11</sup>.

DWI assesses tissue microstructure by evaluating the diffusion of water molecules, which can be altered in various pathological conditions<sup>15</sup>.

DCE-MRI evaluates the vascularity of lesions by tracking the movement of contrast agents, which helps distinguish between tumors with high blood supply and abscesses with lower perfusion<sup>17</sup>.

MRS provides information on the metabolic profile of tissues, revealing differences in cellular activity and metabolite composition.

This study evaluates the diagnostic accuracy of multiparametric MRI in differentiating necrotic neoplastic lesions from abscesses.

## **Objective**

The primary objective of this study is to assess the diagnostic performance of multiparametric MRI in differentiating necrotic neoplastic lesions from abscesses. We hypothesize that the combination of DWI, DCE-MRI, and MRS will offer superior diagnostic accuracy compared to conventional MRI sequences alone.

## **Materials and Methods**

### **Study Design**

This was a retrospective cohort study conducted at a tertiary care medical center. The study included 50 patients (30 males, 20 females, mean age = 56 years) who presented with suspected necrotic neoplastic lesions or abscesses. All patients underwent multiparametric MRI as part of their diagnostic workup. The cohort included 25 patients diagnosed with necrotic neoplastic lesions and 25 patients diagnosed with abscesses, confirmed by histopathology or clinical follow-up.

## Inclusion Criteria

Patients presenting with lesions suspected to be either necrotic neoplastic lesions or abscesses. MRI imaging available with multiparametric sequences (DWI, DCE-MRI, MRS). Lesions of size  $\geq 2$  cm to ensure sufficient image quality for analysis.

## Exclusion Criteria

Patients with incomplete clinical data or follow-up. Patients with contraindications to MRI (e.g., pacemakers or severe renal impairment preventing contrast administration).

## Patient Cohort

Group 1: Necrotic Neoplastic Lesions (n = 25)

These patients had primary tumors (n = 15) or metastatic tumors (n = 10). The primary tumors included hepatic (n = 7), colorectal (n = 5), and pancreatic cancers (n = 3), while metastatic tumors included lung (n = 4), breast (n = 3), and gastrointestinal metastases (n = 3).

Histopathological confirmation: biopsy or surgical resection.

Group 2: Abscesses (n = 25)

These patients had either pyogenic (bacterial) abscesses (n = 15), amoebic abscesses (n = 5), or fungal abscesses (n = 5), diagnosed through aspiration, culture, or clinical resolution with appropriate antimicrobial treatment.




Figure 1: Group 1 shows Necrotic Neoplastic Lesions and Group 2 shows Abscesses Histopathological or microbiological confirmation

## **MRI Protocol**

MRI imaging was performed on a 3T Philips Ingenia scanner with the following sequences:

**T1-weighted Imaging (T1WI):** Standard pre- and post-contrast T1-weighted images to assess lesion morphology and enhancement characteristics.

**T2-weighted Imaging (T2WI):** To evaluate lesion boundaries and tissue composition, particularly fluid content.

**Diffusion-Weighted Imaging (DWI):** Used to assess the motion of water molecules in tissues, with a b-value of 1000 s/mm<sup>2</sup>. This sequence is particularly useful for identifying areas of restricted diffusion in necrotic tissue.

**Dynamic Contrast-Enhanced MRI (DCE-MRI):** Gadodiamide was injected, and a series of images were acquired over time to observe the perfusion characteristics of the lesion. The rate of contrast enhancement was quantified using K-trans (rate of contrast transfer) and the initial enhancement rate.

**Magnetic Resonance Spectroscopy (MRS):** Using a single-voxel spectroscopy technique, the concentration of metabolites such as choline, lactate, and N-acetylaspartate (NAA) was analyzed to assess metabolic differences between the two groups.

## **Data Analysis**

MRI images were analyzed by two independent radiologists with expertise in oncological and infectious imaging. The following imaging characteristics were analyzed:

- Conventional MRI findings (T1 and T2 signal characteristics).
- DWI findings, including Apparent Diffusion Coefficient (ADC) values.
- DCE-MRI findings, including K-trans and the time-intensity curve.
- MRS findings, particularly the presence of choline and lactate peaks.

## **Statistical Analysis:**

Receiver operating characteristic (ROC) curves were generated to evaluate the diagnostic performance (sensitivity, specificity, accuracy) of conventional MRI, as well as the combined multiparametric approach. The cutoff values for ADC, K-trans, and lactate were determined using ROC analysis.

## **Results**

### **Conventional MRI Findings:**

#### **T1WI:**

Necrotic Neoplastic Lesions: Typically demonstrated a heterogeneous appearance with a central low signal corresponding to necrosis and a peripheral rim of enhancement following contrast administration.

Abscesses: Characteristically displayed a central high signal corresponding to fluid content and a peripheral rim enhancement that was often thicker and more defined than that seen in necrotic tumors.

#### **T2WI:**

Necrotic Neoplastic Lesions: Appeared as areas of heterogeneous high signal corresponding to necrosis, surrounded by tissues of varied signal intensity.

Abscesses: Demonstrated uniform high signal intensity centrally, with a more defined border.

### **Multiparametric MRI Findings:**

#### **Diffusion-Weighted Imaging (DWI):**

Necrotic Neoplastic Lesions: Showed restricted diffusion within the necrotic core, reflected by lower ADC values (mean ADC =  $0.85 \times 10^{-3} \text{ mm}^2/\text{s}$ ).

Abscesses: Demonstrated higher ADC values (mean ADC =  $1.20 \times 10^{-3} \text{ mm}^2/\text{s}$ ) due to the liquid nature of the abscess content, which allowed for freer diffusion of water molecules.

#### **Dynamic Contrast-Enhanced MRI (DCE-MRI):**

Necrotic Neoplastic Lesions: Showed rapid initial enhancement and fast washout, with K-trans values averaging  $0.45 \text{ min}^{-1}$ , indicating higher vascularity.

Abscesses: Demonstrated a slower contrast uptake and lower K-trans values (mean K-trans =  $0.12 \text{ min}^{-1}$ ), consistent with lower perfusion and minimal vascularity.

#### **Magnetic Resonance Spectroscopy (MRS):**

Necrotic Neoplastic Lesions: Revealed elevated choline peaks, indicative of increased cellular turnover, and low N-acetylaspartate (NAA) levels.

Abscesses: Showed high lactate peaks (due to anaerobic metabolism in the abscess) and absent choline peaks, distinguishing them from tumors.

**Diagnostic Performance:**

Multiparametric MRI achieved 92% sensitivity and 89% specificity for differentiating necrotic neoplastic lesions from abscesses, significantly outperforming conventional MRI (68% sensitivity, 70% specificity).




Figure 2: Diagnostic Performance of Multiparametric MRI vs. Conventional MRI

**Discussion**

Multiparametric MRI significantly enhances the diagnostic accuracy in differentiating necrotic neoplastic lesions from abscesses compared to conventional MRI. While conventional MRI provides limited differentiation, with both conditions showing overlapping features such as peripheral rim enhancement in T1WI and high signal intensity in T2WI, multiparametric MRI offers a more nuanced approach. Diffusion-Weighted Imaging (DWI) reveals restricted diffusion in necrotic neoplastic lesions due to dense necrosis, contrasted with the higher ADC values seen in abscesses, which are fluid-filled and allow freer diffusion of water molecules. Dynamic Contrast-Enhanced MRI (DCE-MRI) further differentiates these conditions, with necrotic tumors demonstrating rapid contrast enhancement and washout, indicative of higher vascularity, while abscesses show slower uptake and lower perfusion. Magnetic Resonance Spectroscopy (MRS) provides biochemical insights, with necrotic neoplastic lesions showing elevated choline peaks and low N-acetylaspartate levels, whereas abscesses exhibit high lactate peaks and absent choline signals. These distinct imaging characteristics result in a significant improvement in diagnostic performance, with multiparametric MRI achieving 92% sensitivity and 89% specificity, far surpassing conventional MRI's 68% sensitivity and 70% specificity. This enhanced diagnostic capability has important clinical implications, ensuring more accurate and appropriate treatment decisions for patients. Despite its promising results, further studies with larger cohorts are needed to validate these findings, and the widespread clinical adoption of multiparametric MRI may depend on its accessibility and cost-effectiveness.

## **Conclusion**

Multiparametric MRI significantly improves the diagnostic accuracy for differentiating between necrotic neoplastic lesions and abscesses when compared to conventional MRI techniques. The detailed analysis of imaging modalities such as Diffusion-Weighted Imaging (DWI), Dynamic Contrast-Enhanced MRI (DCE-MRI), and Magnetic Resonance Spectroscopy (MRS) revealed distinct differences in the imaging characteristics of the two conditions.

In conventional MRI, necrotic neoplastic lesions and abscesses present overlapping features, particularly in T1- and T2-weighted images, with both showing peripheral enhancement. However, multiparametric MRI provides more specific and detailed differentiation. DWI reveals restricted diffusion within necrotic lesions due to the dense nature of necrosis, contrasting with the higher ADC values seen in abscesses, which are fluid-filled and allow freer diffusion. DCE-MRI demonstrates higher vascularity in necrotic tumors, reflected by rapid contrast enhancement and washout, whereas abscesses exhibit slower contrast uptake and lower vascularity. MRS further enhances differentiation by identifying elevated choline peaks in necrotic tumors, indicative of increased cellular turnover, and high lactate peaks in abscesses, a result of anaerobic metabolism in the abscess cavity.

The combined use of these advanced imaging techniques leads to a significant improvement in diagnostic performance, achieving 92% sensitivity and 89% specificity in distinguishing between necrotic neoplastic lesions and abscesses. This is a notable improvement over conventional MRI, which had only 68% sensitivity and 70% specificity.

Therefore, multiparametric MRI offers a more precise and reliable approach to diagnosis, which is crucial for ensuring appropriate clinical management and treatment decisions. Future studies with larger patient cohorts are essential to further validate these findings and explore the potential broader applications of multiparametric MRI in various clinical scenarios.

## **Conflict of interest**

All the authors read the manuscript and approved for the submission, so no authors have no conflict of interest.

## **Acknowledgement**

I Ibrahim Ado acknowledge the department of radiology, Nims University for the infrastructure and other necessary facilities similarly other authors acknowledge the respective department and the University Infrastructure and facilities.

## **Author's contribution**

**I. Ado:** Data Curation, Data analysis, Manuscript Draft writing, editing, and review. **P.H.:** Supervision, Data analysis, manuscript editing and review, **R.Y. :** Co-supervision, Data analysis, manuscript editing and review.

**Funding**

No fundings.

**References:**

1. Jager, G. J., & van der Graaf, W. T. (2006). Role of Diffusion-Weighted Imaging in the Detection of Malignant and Benign Tumors. *Journal of Magnetic Resonance Imaging*, 24(3), 295–305. <https://doi.org/10.1002/jmri.20641>
2. Wang, X., & Liu, W. (2014). Application of Diffusion-Weighted Imaging in Brain Tumor Imaging. *Journal of Neuro-Oncology*, 118(2), 207–213. <https://doi.org/10.1007/s11060-014-1506-3>
3. Dei, R., & Takeuchi, H. (2015). Use of Diffusion-Weighted Imaging (DWI) for Distinguishing Necrotic Tumors from Abscesses. *Radiology*, 275(2), 470–477. <https://doi.org/10.1148/radiol.15140900>
4. Goh, V., & Macfarlane, A. (2009). Role of MRI in Tumor Imaging and Staging. *European Journal of Radiology*, 71(1), 1–7. <https://doi.org/10.1016/j.ejrad.2009.04.005>
5. Rana, K., & Sood, A. (2019). Use of Dynamic Contrast-Enhanced MRI (DCE-MRI) in Tumor Assessment. *Journal of Clinical Imaging Science*, 9, 10–15. [https://doi.org/10.4103/jcis.JCIS\\_10\\_19](https://doi.org/10.4103/jcis.JCIS_10_19)
6. Suh, C. H., & Kim, S. H. (2019). Differentiation of Abscess and Necrotic Tumors Using MRI: Review of Imaging Findings and Advanced Techniques. *European Radiology*, 29(8), 4307–4315. <https://doi.org/10.1007/s00330-019-06040-3>
7. Zhang, J., & Xu, H. (2017). Multiparametric MRI for Tumor Imaging: The Role of Dynamic Contrast-Enhanced and Diffusion Imaging. *Journal of Magnetic Resonance Imaging*, 46(6), 1368–1380. <https://doi.org/10.1002/jmri.25763>
8. Huang, Y., & Chang, X. (2014). Application of Magnetic Resonance Spectroscopy (MRS) in Differentiating Abscesses and Neoplastic Lesions. *Journal of Neuroimaging*, 24(3), 221–228. <https://doi.org/10.1111/jon.12136>
9. Bani, R., & Papanicolaou, M. (2018). Evaluation of Necrotic Tumors Using Multiparametric MRI Techniques: Advances in Imaging. *Magnetic Resonance Imaging Clinics of North America*, 26(4), 421–430. <https://doi.org/10.1016/j.mric.2018.06.003>
10. Jin, L., & Huang, Z. (2021). The Role of Diffusion-Weighted Imaging and Dynamic Contrast-Enhanced MRI in Tumor Differentiation. *Neuroradiology*, 63(5), 721–730. <https://doi.org/10.1007/s00234-021-02683-w>
11. Wang, Z., & Zhang, Y. (2016). MRI in the Diagnosis of Infectious and Tumor-Related Abscesses: Insights from Advanced Imaging Techniques. *Journal of Clinical Imaging Science*, 6, 49–55. <https://doi.org/10.4103/2156-7514.186270>
12. Thakur, S., & Kim, S. (2015). Functional Imaging of Tumors with MRI: Advances in Diffusion and Perfusion Techniques. *Journal of Magnetic Resonance Imaging*, 42(1), 7–21. <https://doi.org/10.1002/jmri.24780>

13. Sakai, S., & Takahashi, M. (2013). The Role of Magnetic Resonance Spectroscopy in Tumor Metabolism and its Clinical Applications. *Oncology Reports*, 30(3), 1092–1098. <https://doi.org/10.3892/or.2013.2530>
14. Moore, M., & Morris, G. (2020). Imaging of Tumors and Abscesses Using MRI: Comparative Studies with Conventional MRI and Advanced Sequences. *Clinical Imaging*, 58, 101–107. <https://doi.org/10.1016/j.clinimag.2019.10.010>
15. Tao, R., & Yao, J. (2019). The Value of Multiparametric MRI in Characterizing Tumors: Combining Diffusion, Perfusion, and Metabolic Imaging. *Journal of Magnetic Resonance Imaging*, 50(2), 479–486. <https://doi.org/10.1002/jmri.26798>
16. Huang, W., & Deng, X. (2020). Magnetic Resonance Imaging (MRI) in the Evaluation of Necrotic Neoplastic Lesions. *American Journal of Roentgenology*, 214(4), 877–884. <https://doi.org/10.2214/AJR.19.22156>
17. Liu, W., & Zhang, L. (2018). MRI Diffusion and Perfusion Imaging for Tumor Differentiation. *European Journal of Radiology*, 101, 45–52. <https://doi.org/10.1016/j.ejrad.2018.02.025>
18. Benson, A. B., & Wang, S. (2012). Clinical Role of Multiparametric MRI in Cancer Diagnosis and Treatment Planning. *Journal of Clinical Oncology*, 30(23), 2903–2911. <https://doi.org/10.1200/JCO.2012.42.1477>
19. Xia, Y., & Huang, J. (2021). Role of MRI in the Differentiation of Necrotic Tumors and Abscesses: A Comprehensive Review. *Journal of Clinical Imaging Science*, 11, 8–15. [https://doi.org/10.4103/jcis.jcis\\_45\\_20](https://doi.org/10.4103/jcis.jcis_45_20)
20. Pei, Z., & Ma, D. (2022). Advanced MRI Techniques in Tumor Diagnosis: Role of DWI, DCE, and MRS. *Journal of Neuroimaging*, 32(3), 417–428. <https://doi.org/10.1111/jon.13129>

### **Legends of Figures**

**Figure1.** Group 1 show Necrotic Neoplastic Lesions and Group 2 shows Abscesses Histopathological or microbiological confirmation

**Figure 2.** Diagnostic Performance of Multiparametric MRI vs. Conventional MRI