

How to Cite:

Rangari, N. T., More, V. S., Chumbhale, D. S., Kadam, J. N., Kalyankar, T. M., & Muley, Y. P. (2022). Development and validation of stability indicating UV-visible spectrophotometric method for simultaneous determination of salbutamol sulphate and ambroxol hydrochloride in liquid dosage form. *International Journal of Health Sciences*, 6(S1), 2744–2758.
<https://doi.org/10.53730/ijhs.v6nS1.5208>

Development and Validation of Stability Indicating UV-Visible Spectrophotometric Method for Simultaneous Determination of Salbutamol Sulphate and Ambroxol Hydrochloride in Liquid Dosage Form

Nalanda T. Rangari

Faculty of Pharmacy, Alard College of Pharmacy, Marunji, Pune, Maharashtra, India

Vishal, S. More

Faculty of Pharmacy, Amrutvahini College of Pharmacy, Sangamner, Maharashtra, India

Deshraj S. Chumbhale

Faculty of Pharmacy, Amrutvahini College of Pharmacy, Sangamner, Maharashtra, India

Jyoti N. Kadam

Faculty of Pharmacy, Alard College of Pharmacy, Marunji, Pune, Maharashtra, India

Tukaram M. Kalyankar

Department of Pharmaceutical Chemistry, School of Pharmacy, Swami Ramanand Teerth Marathwada University, Vishnupuri, Nanded, Maharashtra, India

Yogiraj P. Muley

Department of Pharmaceutical Chemistry, School of Pharmacy, Swami Ramanand Teerth Marathwada University, Vishnupuri, Nanded, Maharashtra, India



Abstract---Purpose: A new, simple, precise and validated UV spectrophotometric method has been developed for simultaneous estimation of salbutamol sulphate and ambroxol hydrochloride in liquid dosage form. **Method:** The method employed was simultaneous equation method which involves solving simultaneous equations based on measurement of absorbance at two wavelengths 224 nm and 244 nm, the λ_{max} of salbutamol sulphate and ambroxol

hydrochloride, respectively. Specificity of the method was determined by subjecting the drugs to various stress conditions like acid, alkali, thermal and photolytic degradation. *Results:* The linearity was obtained in the concentration range of 0.1-0.6 µg/mL for salbutamol sulphate and 1.5-09 µg/mL for ambroxol hydrochloride. The proposed method was effectively applied to liquid dosage form for estimation of both drugs. The accuracy and reproducibility results are close to 100% with ≤2% RSD. The results of proposed method have been validated as per ICH guidelines. *Conclusion:* A novel, precise and accurate stability indicating spectrophotometric method has been developed for the estimation of salbutamol sulphate and ambroxol hydrochloride in pharmaceutical syrup formulation.

Keywords---Salbutamol sulphate, Ambroxol hydrochloride, Simultaneous equation, Stability indicating, Validation, Spectrophotometric method.

Introduction

Salbutamol sulphate is a chemically bis [(1RS)-2-[(1, 1-dimethylethyl) amino]-1-[4-hydroxy-3 (hydroxymethyl) phenyl] ethanol] sulphate (Fig. 1), it is beta-adrenoceptor agonist used for the relief of broncho-spasm in conditions such as asthma and COPD (chronic obstructive pulmonary disease)¹⁻⁴. It stimulates beta (2)-adrenergic receptors located in the lungs, which ultimately results in relaxation of bronchial smooth muscles⁵. Literature survey reveals that salbutamol sulphate in combination with other drugs has been estimated by UV spectrophotometric methods⁶⁻¹⁰, HPTLC¹¹, RP-HPLC¹², TLC method¹³. Ambroxol hydrochloride is a chemically trans-4-((2-amino-3,5-dibromobenzyl) amino) cyclohexanol hydrochloride (Fig. 2). Ambroxol hydrochloride reduces bronchial hyper-reactivity and acts as a mucolytic and cough suppressant. Both salbutamol sulphate and ambroxol hydrochloride are official in Indian pharmacopoeia. Combination of salbutamol sulphate and ambroxol hydrochloride is used for the treatment of asthma and bronchitis¹⁴⁻¹⁶. Literature survey also reveals UV spectrophotometric¹⁷⁻²⁰, RP-HPLC²¹⁻²⁵, HPTLC²⁶ and LC-MS/MS²⁷ methods are reported for estimation of ambroxol hydrochloride with other drugs in combination. Only UV Spectrophotometric²⁸, RP-HPLC²⁹ and HPTLC³⁰ methods have been found to be reported for the simultaneous estimation of salbutamol sulphate and ambroxol hydrochloride in combination for solid dosage form and no any stability indicating UV spectrophotometric method reported yet for liquid dosage form of same drug combination. Therefore, the objective of the present work is to develop new stability indicating UV spectrophotometric method for estimation of salbutamol sulphate and ambroxol hydrochloride in liquid dosage form with good accuracy, simplicity, precision and reproducibility over other spectrophotometric methods and which can be used for routine analysis methods for simultaneous determination of salbutamol sulphate and ambroxol hydrochloride. The proposed methods were optimized and validated as per ICH guidelines³¹.

Fig. 1: Structure of Salbutamol sulphate**Fig. 2: Structure of Ambroxol hydrochloride**

Materials and Methods

The instrument used in the present study was UV- spectrophotometer UV-1800 (Shimadzu, Japan) with spectral bandwidth of 2 nm and 10 mm matched quartz cells over the range of 200-400 nm. All weighing was done on digital weighing balance (Anamed, AA-2200).

Reagents and Chemicals

Sample of salbutamol sulphate and ambroxol hydrochloride were obtained as a gift sample from Orex Pharma Pvt. Ltd., Thane, India and Erica Pharma Pvt. Ltd., Mumbai, India respectively. Marketed formulation Ambrodil-S syrup containing salbutamol sulphate 01 mg and Ambroxol hydrochloride 15mg for each 5 mL of syrup was purchased from local market. All solvents used were of AR grade.

Preparation of Stock Standard Solutions

Standard stock solutions of both salbutamol sulphate and ambroxol hydrochloride were prepared by dissolving 10 mg of salbutamol sulphate and 10 mg of ambroxol hydrochloride separately in 10 mL of 0.1N HCL in 100 mL volumetric flasks. Final volume was made up to 100 mL with 0.1N HCL to get working standard solution of each containing 100 µg/mL of both salbutamol sulphate and ambroxol hydrochloride.

Determination of Absorption Maxima

By appropriate dilution of standard stock solutions of salbutamol sulphate and ambroxol hydrochloride with 0.1N HCL, solutions containing 10 µg/mL of salbutamol sulphate and 10 µg/mL of ambroxol hydrochloride were scanned separately in the range of 200-400 nm. Wavelength of absorption maxima's was determined for both drugs. Salbutamol sulphate showed maximum absorbance at 224 nm and ambroxol hydrochloride at 244 nm.

Development of Simultaneous Equation

From the stock solution, working standard solution of drugs was prepared by appropriate dilution and was scanned from 400nm to 200nm. Two wavelengths were selected for this method i.e. 224 nm and 244 nm that are absorption maximas of salbutamol sulphate and ambroxol hydrochloride respectively in 0.1N HCL. Series of dilution were prepared from standard solutions of salbutamol sulphate and ambroxol hydrochloride. The linearity was observed in the concentration range of 0.1-0.6 µg/mL for salbutamol sulphate and 1.5- 09 µg/mL for AMB. The absorbance values were measured at the selected wavelengths and absorptivities for both the drugs at both wavelengths were determined. The calibration curves for salbutamol sulphate and ambroxol hydrochloride were plotted in the concentration range of 0.1-0.6 µg/mL and 1.5-09 µg/mL. The concentrations of drugs in sample solution were determined by using the formula 3 and 4,

$$A_1 = ax_1Cx + ay_1Cy \dots\dots\dots(1)$$

$$A_2 = ay_1Cx + ay_2Cy \dots\dots\dots(2)$$

For the measurements in 1 cm cells, b = 1

Rearranging above equations,

$$C_x = \frac{A_2ay_1 - A_1ay_2}{ax_2ay_1 - ax_1ay_2} \dots\dots\dots(3)$$

$$C_y = \frac{A_1ax_2 - A_2ax_1}{ax_2ay_1 - ax_1ay_2} \dots\dots\dots(4)$$

Where, A_1 and A_2 = Absorbance of sample at λ_1 and λ_2


C_x and C_y = Concentrations of salbutamol and ambroxol in sample matrix.

ax_1 and ax_2 = Absorptivities of salbutamol sulphate at λ_1 and λ_2

ay_1 and ay_2 = Absorptivities of ambroxol hydrochloride at λ_1 and λ_2

By solving the two simultaneous equations, the concentrations of salbutamol sulphate and ambroxol hydrochloride in sample solutions were obtained. Overlain spectrum of salbutamol and ambroxol by UV Spectrophotometry is shown in (Fig.3)

Fig. 3: Overlain spectrum of salbutamol and ambroxol by UV Spectrophotometry



Analysis of marketed formulation

Accurately 5 mL of ambrodil-S syrup (Label claim: Each 5mL of syrup contains salbutamol sulphate 01 mg and ambroxol hydrochloride 15 mg) was measured and taken into a 100mL volumetric flask. It is then dissolved in 0.1N HCL and is made up to the mark. The solution is then filtered through Whatman filter paper no. 41. The concentrations of the two drugs in the sample solutions were determined by simultaneous equation (Table 1).

Table 1: Analysis of marketed syrup formulation

* Indicates average of six determinations

Sr. no.	Label claim (mg/5 ml)		Amount of drug found (mg/5 ml)		% Label Claim	
	SAL	ABX	SAL	ABX	SAL	ABX
1	01	15	1.01	15.04	101.34	100.32
2	01	15	1.00	15.15	100.22	101.02
3	01	15	0.99	14.99	99.82	99.98
4	01	15	0.99	14.98	99.97	99.91
5	01	15	0.98	14.79	98.93	98.65
6	01	15	1.00	15.24	100.25	101.65
Mean*					100.08	100.25
SD					0.779	1.030
% RSD					0.778	1.027

Method Validation

Validation of proposed method was done as per ICH guidelines³¹ by studying the following method validation parameters.

Linearity:

As per ICH guidelines the linearity of an analytical procedure is its ability to obtain test results which are directly proportional to the concentration (amount) of analyte in the sample³¹. The linearity of measurement was evaluated by analyzing different concentrations of the standard solution of salbutamol sulphate and ambroxol hydrochloride. For linearity study, stock solutions were subsequently diluted with 0.1 N HCL to get concentrations from 0.1-0.6 µg/ml and 1.5-09 µg/ml of salbutamol sulphate and ambroxol hydrochloride respectively. Then the absorbance of these diluted solutions were measured at 224 nm (λ_1) for salbutamol sulphate and 244 nm (λ_2) for ambroxol hydrochloride by double beam UV spectrophotometer against a blank of 0.1 N HCL. Average of six replicates readings was taken and tabulated (Table 2-4). For both the methods, the Beer law was obeyed in the concentration range 0.1-0.6 µg/mL and 1.5-09 µg/mL for salbutamol sulphate and ambroxol hydrochloride respectively. Calibration curve of salbutamol sulphate and ambroxol hydrochloride is shown in (Fig. 4) and (Fig. 5) respectively.

Fig. 4: Calibration curve of salbutamol sulphate by UV Spectrophotometry




Fig. 5: Calibration curve of ambroxol hydrochloride by UV Spectrophotometry




Table 2: Linearity study data of salbutamol sulphate

Sr. No.	Conc. (µg/ml)	Absorbance at 224 nm
1	0.1	0.022
2	0.2	0.043
3	0.3	0.064
4	0.4	0.081
5	0.5	0.106
6	0.6	0.124

Table 3: Linearity study data of ambroxol hydrochloride

Sr. No.	Conc. (µg/ml)	Absorbance at 244 nm
1	1.5	0.039
2	3.0	0.078
3	4.5	0.116

4	6.0	0.148
5	7.5	0.186
6	9.0	0.218

Table 4: Optical characteristics and other parameters

Parameters	SAL	ABX
Absorption maxima (nm)	224	244
Linearity range (µg/ml)	0.1-0.6	1.5-9.0
Limit of detection (µg/ml)	0.015	0.135
Limit of quantitation (µg/ml)	0.048	0.411
Slope (<i>m</i>)	0.204	0.023
Intercept (<i>c</i>)	0.001	0.005
Regression coefficient (R ²)	0.998	0.998

Repeatability (Intra-assay precision):

To check the degree of repeatability of the methods, suitable statistical evaluation was carried out. Six samples of the syrup formulation were analyzed for the repeatability study. The standard deviation and coefficient of variance was calculated (Table 5).

Table 5: Repeatability Data

No. of Sample	% Recovery	
	SAL	ABX
1	99.24	99.15
2	99.24	99.15
3	101.7	101.6
4	99.24	99.15
5	99.24	101.6
6	99.24	99.15

Mean*	100	100
SD *	1.312	1.841
%RSD*	1.312	0.841

* Indicates average of six determination

Precision:

The precision of an analytical procedure expresses the closeness of agreement (degree of scatter) between a series of measurements obtained from multiple sampling of the same homogeneous sample under the prescribed conditions³¹. The reproducibility of the proposed methods was determined by performing tablet assay at different time intervals on same day (Intraday precision) and on three different days (Inter day precision) (Table 6).

Table 6: Precision data of marketed formulation

Samples	Intraday		Interday	
	SAL	ABX	SAL	ABX
1	98.80	98.31	98.41	99.15
2	101.19	100.00	98.80	100.84
3	98.80	101.68	99.60	99.71
Mean*	99.59	99.99	99.60	99.90
SD*	1.379	1.685	0.606	0.860
%RSD*	1.385	1.685	0.613	0.861

* Indicates average of three determinations

Accuracy (Recovery studies):

The accuracy of an analytical procedure expresses the closeness of agreement between the value which is accepted either as a conventional true value or an accepted reference value and the value found³¹. To ascertain the accuracy of proposed methods, recovery studies were carried out by standard addition method at three different levels (80%, 100% and 120%). Percent recovery for salbutamol sulphate and ambroxol hydrochloride, by both the methods, was calculated (Table 7-8).

Table 7: Recovery study data for method validation of Salbutamol and Ambroxol

Level of Recovery	Amount present (mg)		Added concentration (mg)		Amount recovered (mg)		% Recovery	
	SAL	ABX	SAL	ABX	SAL	ABX	SAL	ABX
80%	01	15	0.8	12	1.772	26.88	98.47	99.59
	01	15	0.8	12	1.793	26.88	99.66	99.59
	01	15	0.8	12	1.764	26.66	98.04	98.75
100%	01	15	01	15	1.997	29.58	99.85	98.63
	01	15	01	15	1.968	29.79	98.42	99.31
	01	15	01	15	1.996	29.79	99.81	99.31
120%	01	15	1.2	17			100.5	
					2.212	32.53	7	101.68
	01	15	1.2	17			101.3	
					2.230	32.43	8	101.36
	01	15	1.2	17			100.5	
					2.212	31.98	7	99.94

Table 8: Statistical validation of recovery study data

Level of Recovery	% Mean recovery*		SD		% RSD	
	SAL	ABX	SAL	ABX	SAL	ABX
80%	98.72	99.31	0.839	0.484	0.850	0.488
100%	99.36	99.08	0.814	0.392	0.819	0.396
120%	100.84	100.99	0.467	0.926	0.463	0.917

* Indicates average of three determinations

LOD and LOQ:

According to ICH guidelines the detection limit of an individual analytical procedure is the lowest amount of analyte in a sample which can be detected but not necessarily quantitated as an exact value. Limit of detection can be calculated using following equation.

$$LOD = 3.3 \times N/S$$

Where, N is the standard deviation of the peak areas of the drug and S is the slope of the corresponding calibration curve.

The quantitation limit of an individual analytical procedure is the lowest amount of analyte in a sample which can be quantitatively determined with suitable precision and accuracy. Limit of quantification can be calculated using following equation.

$$LOQ = 10 \times N/S$$

Where, N is the standard deviation of the peak areas of the drug and S is the slope of the corresponding calibration curve.

Forced degradation study

Specificity of the method was determined by calculating percent amount of possible degradation products produced during the forced degradation study. The forced degradation was carried out with 0.1 NHCL, 0.1 NNaOH, thermal (80 °C) and UV photolysis (365 nm) as per ICH guidelines³²⁻³⁴.

Results

The absorbance of the salbutamol sulphate and ambroxol hydrochloride was measured at 224 nm and 244 nm respectively and calibration curves were plotted as concentrations versus absorbance. The Relative Standard Deviation (RSD) for intra-day analysis of salbutamol sulphate and ambroxol hydrochloride was found to be 1.385 and 1.685 respectively. RSD for Inter-day analysis of salbutamol sulphate and ambroxol hydrochloride was found to be 0.613 and 0.861 respectively. The accuracy and reproducibility is evident from the data as results are close to 100 % and the value of standard deviation and % R.S.D. were found to be < 2 %; shows the high precision of the method. The summary of validated parameters is shown in (Table 09). The proposed method is simple, economical, rapid, precise and accurate. Hence it can be used for routine analysis of salbutamol sulphate and ambroxol hydrochloride in pharmaceutical formulation. The proposed method was found to be specific as there is no interference from other excipients. The LOD for salbutamol sulphate and ambroxol hydrochloride was found to be 0.015 µg/mL and 0.135µg/mL respectively. The LOQ for salbutamol sulphate and ambroxol hydrochloride was found to be 0.048µg/mL and 0.411µg/mL respectively. Marketed brand of syrup (Ambrodil-S) was analyzed, the amounts of salbutamol sulphate and ambroxol hydrochloride determined by proposed method were found to be 100.08 % and 100.25 % respectively. The stress degradation studies showed that salbutamol sulphate undergoes degradation in acidic, alkaline, photolytic and thermal condition and ambroxol hydrochloride undergoes degradation in acidic, alkaline, photolytic and thermal condition respectively. The results for forced degradation are shown in (Table 10).

Table 9: Summary of validation parameters.

Parameters	Salbutamol sulphate	Ambroxol hydrochloride
Linearity Range (µg/mL)	0.1-0.6	1.5-9.0
Correlation coefficient (r ²)	0.998	0.998
Precision (%RSD)		
Intraday	1.385	1.685
Interday	0.613	0.861
Accuracy (%)		
80% ± RSD	98.72 ± 0.850	99.31 ± 0.488
100% ± RSD	99.36 ± 0.819	99.08 ± 0.396

120% $\pm$ RSD	100.84 $\pm$ 0.463	100.99 $\pm$ 0.917
Repeatability (%RSD)	1.312	1.841
LOD ($\mu\text{g/mL}$)	0.015	0.135
LOQ ($\mu\text{g/mL}$)	0.048	0.411
Solution stability	Stable for 24 hrs.	Stable for 24 hrs.

Table 10: Summary of results of stress degradation studies

Sr. no.	Condition	% Degradation		% Assay	
		SAL	ABX	SAL	ABX
1	Acid hydrolysis (0.1 N HCL, 80°C, 6 hrs.)	38.09	47.61	61.91	52.95
2	Base hydrolysis (0.1 N NaOH, 80°C, 2 hrs.)	20.23	19.32	79.77	80.68
3	Photolytic degradation (UV Radiation, 3 hrs.)	38.09	45.65	61.91	54.35
4	Thermal degradation (80°C, 5 hrs)	17.85	16.52	82.15	83.48

Discussion

In simultaneous equation method, the primary requirement for developing a method for analysis is that the entire spectra should follow the Beer's law at all the wavelength, which was fulfilled in case of both these drugs. The wavelengths used for the analysis of the drugs were 224 nm (λ -max of salbutamol sulphate) and 244 nm (λ -max of ambroxol hydrochloride) at which the calibration curves were prepared for both the drugs. In 0.1 N HCL, salbutamol sulphate and ambroxol hydrochloride obeyed linearity in the concentration range of 0.1-0.6 $\mu\text{g/mL}$ and 1.5-09 $\mu\text{g/mL}$ respectively at their respective λ max with correlation coefficient ($r^2 > 0.99$) in both the case. In proposed method precision was studied as repeatability (%RSD<2) and inter and intra-day variations (%RSD<2) for both drugs; shows the high precision of the method. The accuracy of method was determined by calculating mean percentage recovery. It was determined at 80, 100 and 120 % level and data tabulated. The developed method for estimation of salbutamol sulphate and ambroxol hydrochloride in liquid dosage form was found to be simple, accurate, reproducible, sensitive and economic. For projected method we used easily available and cheap solvent like 0.1 NHCL (AR grade). The proposed method of simultaneous estimation not required any expensive and satisfactory apparatus in contrast to reported chromatographic and hyphenated techniques. So it shows projected method is simple, economic and rapid for estimation of salbutamol sulphate and ambroxol hydrochloride in combined liquid

dosage forms. Hence the developed method for estimation of salbutamol sulphate and ambroxol hydrochloride can be constructive in the routine analysis.

Conclusion

The developed and validated stability indicating UV spectrophotometric method reported here is rapid, simple, accurate, sensitive and specific in accordance to the ICH guidelines showing linearity, accuracy, precision, selectivity, stability and system suitability for the routine determination of salbutamol sulphate and ambroxol hydrochloride. The developed method was successfully used for quantitative estimation and analysis of salbutamol sulphate and ambroxol hydrochloride in combined liquid dosage form. Stress degradation results of show that the method is selective and stability indicating. Thus the reported method is of substantial importance and has great industrial applicability for quality control and analysis of salbutamol sulphate and ambroxol hydrochloride in liquid dosage forms. By observing validation parameter and statistical data, the proposed method was found to be satisfactory over other reported spectrophotometric and chromatographic methods.

Acknowledgements

The authors are thankful to Director, School of Pharmacy, S. R. T. M. University, Nanded, for encouragement and availing of the necessary facilities during the course of investigation. Authors are also gratified to Orex Pharma Pvt. Ltd., Thane, India and Erica Pharma. Pvt. Ltd., Mumbai, India for providing gift sample of salbutamol sulphate and ambroxol hydrochloride respectively.

References

1. *Indian Pharmacopoeia*. 3rd ed. New Delhi: The controller of publication; 2007. P. 1687.
2. John HB, John MB. *Wilson and Gisvold's Textbook "Organic Medicinal and Pharmaceutical chemistry"*. London, UK: Wolters Kluwer; 2004. P. 96-99.
3. Barnes PJ, Page CP. *Pharmacology and therapeutics of asthma and COPD*. Germany: Springer-Verlag Berlin Heidelberg; 2004. P. 15.
4. The Merk Index. *An Encyclopedia of Chemicals, Drugs and Biologicals*. 14th ed. Merk laboratories; 2006. P. 216.
5. Drug Bank: Salbutamol sulphate (DB01001), <http://www.drugbank.ca/drugs/DB01001.pdf>. 2014.
6. Dave HN, Mashru RC, Thakkar AR. Simultaneous determination of salbutamol sulphate, bromhexine hydrochloride and etofylline in pharmaceutical formulations with the use of four rapid derivative spectrophotometric methods. *Anal ChimActa* 2007; 597: 113-120.
7. Mukherji G, Aggarwal N. Quantitative estimation of salbutamol sulphate by derivative UV spectroscopy in the presence of albumin. *Int J Pharm* 1992; 86: 153-158.
8. Mukherji G, Aggarwal N. Derivative UV-spectroscopic determination of salbutamol sulphate in the presence of gelatin. *Int J Pharm* 1991; 71: 187-191.

9. Parimoo P, Umapathi P, Ilango K. Simultaneous quantitative determination of salbutamol sulphate sulfate and bromhexine hydrochloride in drug preparations by difference spectrophotometry. *Int J Pharm* 1993; 100: 227-231.
10. Patel PA, Dole MN, Sawant SD, Shedpure PS. Simultaneous determination of salbutamol sulphate and ambroxol in fixed dose combination by spectrophotometry. *Int J Pharma Sci and Research* 2011; 2(5): 1225-1230.
11. Colthup PV, Dallas FA, Saynor DA, Carey PF, Skidmore LF, Martin LE. Determination of Salbutamol sulphate in human plasma and urine by high-performance thin-layer chromatography. *J Chromatogr.* 1985; 345: 111-118.
12. Jacobson GA, Peterson GM. High-performance liquid chromatographic assay for the simultaneous determination of ipratropium bromide, fenoterol, salbutamol sulphate and terbutaline in nebulizer solution. *J Pharm & Biomed Anal* 1994; 12: 825-832.
13. Dave HN, Mashru RC, Patel AK. Thin Layer Chromatography Method for the Determination of Ternary Mixture Containing Salbutamol Sulphate, Bromhexine Hydrochloride and Etofylline. *J Pharm Sci & Res* 2010; 2(2): 143-148.
14. *Indian Pharmacopoeia*, 3rd ed. New Delhi: The controller of publication; 2007. P. 143,250, 701.
15. The Merk Index. An Encyclopedia of Chemicals, Drugs and Biologicals. 14th ed. Merk laboratories; 2006. P. 385.
16. Barnes PJ, Page CP. Pharmacology and therapeutics of asthma and COPD. Germany: Springer-Verlag Berlin Heidelberg; 2004. P. 217-218.
17. Gowekar NM, Pande VV, Kasture AV, Tekade AR, Chandorkar JG. Spectrophotometric estimation of ambroxol and cetirizine hydrochloride from tablet dosage form. *Pak J Pharm Sci* 2007; 20(3): 250-251.
18. Bhatia NM, Ganbavale SK, Bhatia MS, More HN, Kokil SU. RP-HPLC and Spectrophotometric estimation of ambroxol hydrochloride and cetirizine hydrochloride in combined dosage form. *Ind J Pharm Sci* 2008; 70: 603-608.
19. Makarand A, Bonde CG. Development and validation of simultaneous UV spectrophotometric method for the determination of levofloxacin and ambroxol in tablets. *Int J Chem Tech Research* 2009; 1: 873-888.
20. Hadad GM, Gindy AE, Waleed MM. HPLC and chemometrics-assisted UV-spectroscopy methods for the simultaneous determination of ambroxol and doxycycline in capsule. *SpectrochimActa Part A* 2008; 70: 655-663.
21. Dinçer Z, Basan H, Goger NG. Quantitative determination of ambroxol in tablets by derivative UV spectrophotometric method and HPLC. *J Pharm & Biomed Anal* 2003; 1: 31(5): 867-872.
22. Nobilis M, Pastera J, Svoboda D and Kvstina J. High-Performance Liquid Chromatographic determination of ambroxol in human plasma. *J Chromatogr* 1992; 581: 251-255.
23. Meiling Qi, Wang P, Cong R, Yang J. Simultaneous determination of roxithromycin and ambroxol hydrochloride in a new tablet formulation by liquid chromatography. *J Pharm & Biomed Anal* 2004; 35:1287-1291.
24. Nagappan KV, Meyyanathan SN, Raja RB, Reddy S, Jeyaprakash MR, Birajdar AS. A RP-HPLC Method for Simultaneous Estimation of Ambroxol

- Hydrochloride and Loratidine in Pharmaceutical Formulation. *Research J Pharm and Tech* 2008; 4: 366-369.
25. Maithani M, Raturi R, Vertika Gautam, Simultaneous estimation of ambroxol hydrochloride and cetirizine hydrochloride in tablet dosage form by RP-HPLC method. *Int J Comprehensive Pharm* 2010; 1: 1-3.
 26. Jain PS. Stability-indicating HPTLC determination of ambroxol hydrochloride in bulk drug and pharmaceutical dosage form. *J Chromatogr Sci* 2010; 48(1): 45-48.
 27. Hohyun K, Jeong-Yeon Y, Sang BH, Hee JL, Kyung RL. Determination of ambroxol in human plasma using LC-MS/MS. *J Pharm & Biomed Anal* 2003; 32: 209-216.
 28. Patel PA, Dole MN, Shedpure PS, Sawant SD. Spectrophotometric simultaneous estimation of salbutamol sulphate and ambroxol in bulk and formulation. *Asian Journal of Pharmaceutical and Clinical Research* 2011; 4 (3): 42-45.
 29. Chitlange SS, Dhole AG, Pandkar SV, Wankhede SB. Development and validation of RP-HPLC method for the simultaneous estimation of salbutamol sulphate and ambroxol hydrochloride in tablet dosage form. *Inventi Rapid: Pharm Ana and QualAssur* 2011; 2:131.
 30. Deosarkar AV, Deshpande SD, Walode SG, Tuljapure DS, Tekale SG, Waghmode VM. Simultaneous quantification of salbutamol sulphate and ambroxol hydrochloride by rp-HPLC and HPTLC in bulk drug and dosage form. *International Journal of Pharmacy and Pharmaceutical Sciences* 2012; 4(4): 307-311.
 31. International Conference on Harmonization (ICH), Draft guidelines on validation of Analytical Procedure, Definition and Terminology. Geneva: Federal Register; 1995. p. 11260-62.
 32. International Conference on Harmonization (ICH), Draft guidelines on Stability Testing of New Drug Substances and Products. Geneva: Federal Register; 1999.
 33. Guidance for Industry (Q1B Photostability Testing of new drug Substances and Products). November 1996.
 34. Guidance for Industry (Stability testing of drug substances and drug products). June 1998.