KNOWLEDGE ON ROLE OF BREAST CRAWL IN MATERNAL HEALTH AND WELL-BEING AMONG NURSES

Christena.P*, Elizebeth Rani.V**, Sandhiya.T***

*Lecturer, **Reader, *** Staff Nurse Srinivasan Nursing College, Trichy M.A.Chidambaram College of Nursing, Chennai VHS hospital

Abstract

Introduction: Uninterrupted Breast crawl technique is important just after the delivery, may promote breastfeeding and prevent postpartum Introduction: Office of the safe group of 25-40 years were selected through non-probability corrections of the safe group of 25-40 years were selected through non-probability corrections. hemorrnage. In the age group of 25-40 years were selected through non-probability convenience sampling technique. Results: The findings of the wellbeling. 30 states the state of the research project revealed that 70 % nurses had adequate knowledge on role of breast crawl in maternal health and well-being and 30% nurses had inadequate research project revealed that 70 % nurses had inadequate research project revealed that 70 % nurses had inadequate research project revealed that 70 % nurses had adequate knowledge on role of breast crawl in maternal health and well-being and 30% nurses had inadequate knowledge on role of breast crawl in maternal health and well-being.

Key words: breast crawl technique, breast crawl in maternal health and well being

Introduction

Breast crawl technique in the third stage also benefits the mother; massage of the breast by the baby and subsequent suckling induce a large oxytocin surge from the mother's pituitary gland into her bloodstream. Close emotional interaction coupled with cutaneous visual and auditory stimuli from the baby during Crawl also help oxytocin release. The helps to contract the uterus, explaing the placenta and closing off many blood vessels in the uterus, thus reducing blood loss and preventing anemia. Pressure of the infant's feet on the abdomen may also assist in expelling the placenta (Klaus and Kennel, 2001).

The study shown that baby in Breast Crawl position, the blood level of oxytocin soon after delivery was elevated significantly compared with that postpartum. The level returned to its pre-partum value 1 hour after delivery. The peak oxytocin level was seen 15 minutes after delivery with expulsion of placenta. Most mothers had several peaks of oxytocin occurring up to 1 hour after delivery (Nissen et al, 1995).

Mothers and babies should be together, in breast crawl technique immediately after birth. The baby is happier, the baby's temperature is more stable, the baby's heart and breathing rates are more stable and the baby's blood sugar is more elevated. Breast crawl technique immediately after birth allows the baby to be colonized by the same bacteria as the mother and is thought to be important in the prevention of allergic diseases. When a baby is put into an incubator, his skin and gut are often colonized by bacteria different from his mother's.(Jack Newman, 2005).

The investigator had the chance to see the video when she visited the UNICEF website. She is attracted by the video and practiced the technique in her clinical posting. Therefore, she takes this study to educate and motivate the hospital staffs and other workers for practicing this initiative which will be in turn contributing to healthy children and thereby healthy future citizens of our country. Like all other gifts of nature, this gift comes free of cost. However, the health and nutrition benefits of Breast Crawl will

save millions of life and also will save billions in terms of health cost. They will create a generation which will reach the highest human potential of growth and development.

Statement of the problem

A study to assess the knowledge on role of breast crawls in maternal health and well-being among nurses at selected hospital, Chennai.

Objectives of the study

- 1. To assess the knowledge on role of breast crawl in maternal health and well-being among nurses.
- 2. To associate the knowledge on role of breast crawl in maternal health and wellbeing with demographic variables.

Null Hypotheses

H₀₁ - There is no association between knowledge on role of breast crawl with demographic variables.

Materials and Methods

The research design adopted for this study was Non experimental descriptive research design. 30 staff nurses in the age group of 25-40

years were selected through non-probable convenience sampling technique. The two sections. Section A convenience consisted two sections. Section A consisted two sections as age of the two sections are the two sections as age of the two sections. demographic variables such as age, education status, area of work, working experience years, source of information about role of bre crawl. Section B included structure questionnaire to assess the knowledge on breaching data crawl. The demographic data were collected through Interview guide, structured questionnaire were administered to assess the knowledge.

Descriptive statistics as frequency and percentage distribution, inferential statistic such as chi-square test was used to analyze the

Result& Discussion

The findings related to demographic variables revealed that, Majority (90%) of the sample declong to the age group of 25-35 years. the samples were completed BSc % were working in medical area 87% had less than 5 years of experience, 87% of them had source of information about role of breast crawl through magazines/books.




Fig 1: Percentage distribution of knowledge on role of breast crawl in

TNNMC JOGN Vol. VIII / Issue 2 / Jul - Dec 2020

The above figure shows that 70% of the nurses had adequate knowledge and 30% of the nurses had inadequate knowledge on role of breast crawl in maternal health and well-being.

There was a statistically significant association between level of knowledge with the age at p<0.05 level. There was no statistically significant association between level of knowledge with other demographic variables such as educational status, area of work, working experience in years, source of information about role of breast crawl.

Conclusion

The study concluded that breast feeding is a natural human activity, difficulty is not uncommon. Breast crawl technique is important just after the delivery, the baby sucking reflex is strong and baby is more alert. Putting the baby to the breast as soon as possible after birth helps to avoid many problems. Nurses as health professional, if work in co-ordination as a team to bring forth and promote Breast Crawl initiative into maternity hospitals, it will be beneficial for mother and baby as well as for the entire family and country.

Recommendations

Based on the present study in view, the following recommendations were made,

- A comparative study can be on breast crawl technique in maternal health and well-being in rural and urban areas
- A self-instructional module in local languages can be prepared for educating the mother and their dually members
- The research study can be conducted on a large sample at different settings.

References

- AmandaHenderson. (2014) Promoting the Breast Crawl.http://www.health4mom.org). Retrieved September 2014
- Anne Saxton and Kathleen Fahy (2014). Effects of skin-to-skin contact and breastfeeding at birth on the

- incidence of PPH: A physiologically based theory. Global Medical Discovery 0.1/2.015. Retrieved from www.researchgate.net
- Dr. Sarah J. (2015) Early Skin-to-Skin Contact between Mothers and Newborns-Hormonal Physiology of Childbearing: Evidence and Implications for Women, Babies, and Maternity Care. (childbirthconnection .org). Retrieved 2015
- Elizabeth R Moore. (2012). Early skin-toskin contact for mothers and their healthy newborn infants. (http://www.ncbi.nlm.nih.gov).Retriev ed May 2012
- Healthy Children Project. (2011) The Magical Hour. (http://www. magicalhour.com). Retrieved 2011
- King Edward Memorial Hospital. (2014) Third Stage: Expectant (Physiological) Management Following a Vaginal Birth. (http://www.kemh. health.wa.gov.au).Retrived February 2014
- Ms. Jacintha Lobo. (2014) A quasi experimental study to assess the effectiveness of breast crawl as a diversional therapy for reduction of pain in primipara women, during episiotomy suturing in selected maternity hospitals of a metropolitan city". Retrieved from https://www.researchgate.net 2014.