



Published in final edited form as:

Multivariate Behav Res. 2009 March 1; 44(2): 213–232.

Mediation and moderation: Testing relationships between symptom status, functional health, and quality of life in HIV patients

Ehri Ryu,

Department of Psychology, Boston College

Stephen G. West, and

Department of Psychology, Arizona State University

Karen H. Sousa

College of Nursing, University of Colorado Denver.

Abstract

We extended Wilson and Cleary's (1995) health-related quality of life model to examine the relationships among symptoms status (Symptoms), functional health (Disability), and quality of life (QOL). Using a community sample ($N = 956$) of male HIV positive patients, we tested a mediation model in which the relationship between Symptoms and QOL is partially mediated by Disability. Common and unique ideas from three approaches to examining moderation of effects in mediational models (Edwards & Lambert, 2007; Preacher, Rucker, & Hayes, 2007; MacKinnon, 2008) were used to test whether (a) the direct relationship of Symptoms to QOL and (b) the relationship of Disability to QOL are moderated by age. In the mediation model, both the direct and the indirect (mediated) effects were significant. The direct relationship of Symptoms to QOL was significantly moderated by age, but the relationship of Disability to QOL was not. High Symptoms were associated with lower QOL at all ages, but that this relationship became stronger at older ages. We compare the three approaches and consider their advantages over traditional approaches to combining mediation and moderation.

Keywords

mediation; moderation; direct effect; indirect effect; moderated path analysis; HRQOL

Correspondence: Stephen G. West Department of Psychology Arizona State University PO Box 871104 Tempe AZ 85287-1104 sgwest@asu.edu Phone: 480-965-7339. Correspondence should be directed to Ehri Ryu, Department of Psychology, Boston College, 140 Commonwealth Avenue, Chestnut Hill MA 02467-3807. ryueh@bc.edu..

¹The data were collected from 6 locations corresponding to large geographic areas (communities) within these cities. We calculated the intraclass correlations for Symptoms, Disability, QOL, and Age. The intraclass correlations ranged from 0.04 to 0.08. We used the Complex procedure in Mplus to produce correct standard errors that take into account this dependency in the data. Because of the concerns about participant anonymity, identification numbers for individual clinics were not typically available in the dataset.

²The indirect or mediated effect can be calculated in two ways: (1) difference between total effect (regression of Y on X) and direct effect (regression of Y on X controlling for M), and (2) product of path coefficient from X to M and path coefficient from M to Y . These two methods yield identical estimates of mediated effect when the dependent variable is continuous and ordinary least squares regression is used (MacKinnon, Warsi, & Dwyer, 1995).

³We thank an anonymous reviewer for suggesting this test.

⁴Other statistical programs (e.g., SPSS, SAS, LISREL, EQS) can be used to estimate the models presented here. Edwards and Lambert (2007) provides SPSS syntax in their appendix, SPSS macros are available in Preacher's website (www.quantpsy.org), the CD accompanying MacKinnon (2008) contains computer script and outputs from SPSS, SAS, LISREL, EQS, MPLUS, and CALIS programs.

The management of symptoms is important for the quality of life of patients. Wilson and Cleary (1995) proposed a health-related quality of life (HRQOL) conceptual model that suggests the relationship between symptom status and quality of life is mediated by an individual's functional health status (disability). Parts of this HRQOL conceptual model have been widely applied to different patient populations, including patients with HIV/AIDS (Cunningham, Crystal, Bozzette, & Hays, 2005; Sousa, Holzemer, Henry, Slaughter, 1999; Wilson & Cleary, 1997). In two previous studies, Sousa and Kwok (2005, 2006) found a significant relationship between symptom status and quality of life. It is important to more fully understand the relationships among symptom status, functional health, and quality of life in the HIV population, particularly given the wide range of ages now represented in this population. Determining the effects of age on the relationships between symptom status, functional health, and quality of life could provide beneficial information to develop interventions to support successful aging. We sought to answer four questions: (1) In the HIV/AIDS population, is functional health a mediator between symptom status and quality of life?; (2) Is the relationship between symptom status and quality of life moderated by age?; (3) Is the relationship between functional health and quality of life moderated by age?; and (4) Does age moderate the indirect relationship between symptom status and quality of life that is mediated by functional health?

A secondary goal of this paper is to illustrate the application of new analytic approaches to the study of mediation and moderation. We briefly review mediation, moderation, and traditional approaches to combining mediation and moderation. We then consider three recent approaches (Edwards & Lambert, 2007; MacKinnon, 2008; Preacher, Rucker, & Hayes, 2007) that offer tests of combinations of mediation and moderation in the context of the standard three variable mediational model. These approaches overlap greatly, but each offers some unique features that may be useful in studying different questions involving mediation and moderation. We utilize relevant features from these approaches to test hypotheses about the potential moderation of both direct and indirect effects between symptom status and quality of life in a large sample of HIV/AIDS patients.

Mediation

When mediation is hypothesized, an independent variable (X) is expected to affect an intervening variable (M), which, in turn, is expected to affect a dependent variable (Y). For example, the relationship between neighborhood disadvantage (a single variable characterized by high rates of poverty, crime, and unemployment) and children's externalizing behavior is mediated by the intervening variable of parent-child conflict (Roosa et al., 2005); the relationship between neighborhood disadvantage and children's internalizing behavior is mediated by mothers' perceptions of neighborhood quality such as cleanliness (Deng et al., 2006). In the framework of path analysis, the mediated effect is referred to as an *indirect* effect (Alwin & Hauser, 1975; Bollen, 1987; Fox, 1980). Complete mediation requires that the full effect of the independent variable on the dependent variable be carried by the mediator. Partial mediation recognizes that independent variables may have their own direct effects on the dependent variable that are independent of the mediator. These direct effects may be theoretically important in some contexts and in practice are often larger in magnitude than the mediated effect.

Kenny and his colleagues (Baron & Kenny, 1986; Judd & Kenny, 1981; Kenny, Kashy, & Bolger, 1998) have developed the most widely used procedure for testing for the existence of a mediational effect. Figure 1 illustrates the single mediator model and presents path diagrams illustrating the models and defining the notation below. Their procedure consists of four causal steps that are needed to establish mediation:

Step 1. Show that X and Y are related, i.e., \hat{c} is significant in (1).

$$Y = i_1 + cX + e_1 \quad (1)$$

Step 2. Show that X and M are related, i.e., \hat{a} is significant in (2).

$$M = i_2 + aX + e_2 \quad (2)$$

Step 3. Show that M has an effect on Y over and above X , i.e., \hat{b} is significant in (3).

$$Y = i_3 + c'X + bM + e_3 \quad (3)$$

Step 4. Show that X has no direct effect on Y , i.e., \hat{c}' does not differ from zero in (3).

This final step is a test that the effect is fully carried by the putative mediator (complete mediation).

In contrast, MacKinnon and colleagues (MacKinnon et al., 2002; MacKinnon, 2008) have emphasized focused tests of the indirect $X \rightarrow M \rightarrow Y$ effect. Focused tests provide estimates and tests of significance of the mediated effect, whereas the causal steps procedure only provides a binary conclusion that mediation exists or not. MacKinnon et al. (2002) showed that the causal steps procedure has low statistical power to detect the indirect effect. Correspondingly, Fritz and MacKinnon (2007) showed that exceedingly large sample sizes (e.g., $n > 20,000$) are required to achieve .80 power to detect an indirect effect in which both a and b are small in magnitude and there is no direct effect. Note further that Step 1 represents a test of the total effect, the combination of the direct effect of X on Y and the indirect effect of $X \rightarrow M \rightarrow Y$. Hence, the causal steps approach will fail to detect inconsistent mediation in which the direct and indirect effects are opposite in sign. In addition, given that in practice the magnitude of the direct effect is typically larger than that of the indirect effect, the outcome of this test will often be primarily determined by the magnitude of the direct effect, contrary to the hypothesis of interest. Consequently, Sobel (1982), MacKinnon et al. (2002), MacKinnon, Lockwood, and Williams (2004), and Shrout and Bolger (2002) proposed focused tests of the indirect effect. MacKinnon et al. (2002) provided analytic approaches that are expected to produce proper standard errors given the non-normality of the ab product term, whereas MacKinnon et al. (2004) and Shrout and Bolger (2002) used bootstrapping procedures to achieve this same goal. By focusing on the indirect effect in the testing procedure, these approaches also help refocus attention on the direct effect when it may be of theoretical interest.

Moderation

Moderation occurs when the direction, strength, or both of the relation between an independent and a dependent variable are affected by a third variable, which is termed a *moderator* (Baron & Kenny, 1986; Chaplin, 2007). For example, the efficacy of treatment for depression depends on the patients' attachment insecurity, such that cognitive-behavioral therapy is more effective than interpersonal psychotherapy for patients who are higher in attachment avoidance (McBride, Atkinson, Quilty, & Bagby, 2006). In this example, the effect of therapy on depression reduction is *moderated* by the patients' attachment insecurity. In multiple regression, moderation is typically represented as the linear by linear interaction between the independent variable and the moderator variable (Aiken & West, 1991), using equation (4) below:

$$Y=b_0+b_1X+b_2Z+b_3XZ+e \quad (4)$$

where X is the independent variable and Z is the moderator. The test of the interaction term, b_3 , represents the test of the moderation effect.

Moderation of Indirect and Direct Effects in Mediation Models

Research questions often require examination of moderation of effects in mediational models. As one example, the influence of ethnic cultural norms on cigarette use in Mexican-American youth has been hypothesized to be mediated by self-efficacy. However, the magnitude of the indirect effect decreases as peer smoking increases (Morgan-Lopez, Castro, Chassin, & MacKinnon, 2003). As another example, the relationship between Attention-Deficit / Hyperactivity Disorder (ADHD) and depression is mediated by others' (e.g., teachers and parents) appraisals of social competence. However, the indirect effect holds only in younger children. In other words, the mediated effect of others' appraisals of social competence is moderated by children's age (Ostrander, Crystal, & August, 2006). Researchers have hypothesized a number of different models that combine mediation and moderation. These models have also more recently begun to include hypotheses about moderation of direct effects in mediational models.

Building on initial work by Baron and Kenny (1986) in the causal steps tradition, researchers have historically conducted analyses combining mediation and moderation using one of three general approaches (Edwards & Lambert, 2007). In the *piecemeal approach*, researchers have conducted mediation analysis and moderation analysis separately on the same data set. Mediation is usually tested using the causal steps approach and moderation is tested using analysis of variance or regression analysis. The results of the two sets of analysis are interpreted together. In the *subgroup approach*, the sample is typically dichotomized into subgroups based on the moderator variable (e.g., younger / older children) and a mediational analysis is conducted separately in each subgroup. If the mediation effect differs between subgroups (e.g., mediation effect is significant in one group but not in the other; the magnitude of mediation effect differs between the subgroups), then an inference is made that mediation is moderated. In the *moderated causal steps approach*, mediation analysis is conducted following the causal steps procedure with the product term of independent variable \times moderator added (Baron & Kenny, 1986; Muller, Judd, & Yzerbyt, 2005).

These earlier methods all inherit the shortcomings of the causal steps approach (Baron & Kenny, 1986; Judd & Kenny, 1981) identified above. All of these earlier methods require a significant total X to Y relationship in Step 1, which (a) is often highly dependent on the existence of a direct effect from X to Y and (b) is often not fulfilled under inconsistent mediation. None of the approaches introduced above provide a direct test of indirect effect or provide an estimate of the magnitude of the indirect effect conditioned on the level of the moderator. None of these traditional general methods provides a clear mechanism for modeling which paths (e.g., X to M , M to Y , or X to Y) are affected by the moderator. None of the approaches consider potential moderation of the direct effect. Finally, in the subgroup approach, continuous moderators are dichotomized and moderation is often inferred when an effect is statistically significant in one group, but not in the other, two practices which can be problematic (MacCallum, Zhang, Preacher, & Rucker, 2002; McNemar, 1960).

In response to limitations of previous approaches, Edwards and Lambert (2007), MacKinnon (2008, chapter 10), and Preacher, Rucker, and Hayes (2007) have independently proposed general frameworks for considering moderated effects in mediational models. Each of the

approaches uses a general path analytic framework and develops the regression equations necessary to specify potential models of interest. Consequently, there is considerable overlap among the three approaches. The approaches differ in the models that are emphasized and in subsidiary procedures used in model testing and interpretation.

Edwards and Lambert (2007) argued that prior research and theory might lead researchers to hypothesize that any of the paths in Figure 1 (B) might be moderated or not moderated. Given the three potential paths in the basic mediational model, $X \rightarrow Y$, $X \rightarrow M$, $M \rightarrow Y$, eight ($2 \times 2 \times 2$) potential models may be described, the basic mediational model plus seven additional models that involve moderation of 1, 2, or all 3 paths. Unlike Preacher et al., they explicitly emphasized models that involve the moderation of direct as well as indirect effects. They developed reduced form equations for each of the models that involve the product of the two path coefficients and they recommended the use of bootstrapping methods to construct confidence limits for the product of the two path coefficients. They also extended procedures developed by Aiken and West (1991) to plot and test how indirect and direct effects across levels of the moderator by calculating conditional indirect and direct effects.

Preacher et al. (2007) took a similar general approach to combining moderation and mediation. Preacher et al. (p. 194) considered five models, three of which were identical with models considered in detail by Edwards and Lambert: Model 2, moderation of the a path from the independent variable X to the mediator M by the moderator variable W ; Model 3, moderation of the b path from the mediator M to the dependent variable Y by the moderator variable W ; Model 5, simultaneous moderation of both the X to M and M to Y paths by the common moderator W . In addition, Preacher et al. proposed two models not considered by Edwards and Lambert. Model 4 is a small variant of Model 5 in which W moderates the X to M path and Z moderates the M to Y path. In Model 1, the independent variable X moderates the path from M to Y , a case emphasized by Kraemer, Wilson, Fairburn, and Agras (2002) and discussed earlier by Baron and Kenny (1986). Even though Preacher et al.'s two additional models were not presented in Edwards and Lambert, the general framework of Edwards and Lambert can be easily extended to analyze these models. For the significance tests of the conditional indirect effects, Preacher et al. proposed using bootstrapping methods to construct confidence intervals. They also provided the formulas to obtain asymptotic normal-theory standard errors for key parameters in each of the five models. In addition to the methods presented by Edwards and Lambert for plotting and testing conditional indirect effects, Preacher et al. also proposed an extension of the Johnson and Neyman (1936) procedure that determines the regions of significance over which the conditional indirect effect will be significant.

Finally, MacKinnon (2008) presents a complete volume devoted to mediational analysis. He presents the most general approach (pp. 286-287) using two general algebraic equations that encompass nearly all possible models involving mediation and moderation. All but one of the models proposed by either of Edwards and Lambert or by Preacher et al. can be considered to be special cases of MacKinnon's general approach.

In MacKinnon's (2008) approach, the first equation for predicting the mediator is:

$$M = i_2 + a_1X + a_2Z + a_3XZ + e_2. \quad (5)$$

In this equation the test of a_3 represents the test of the moderation of the X to M path. The second equation for predicting the dependent variable Y is:

$$Y = i_3 + c'_1 X + c'_2 Z + c'_3 XZ + b_1 M + b_2 MZ + hXM + jXMZ + e_3. \quad (6)$$

In this equation, b_2 represents the test of moderation of the M to Y path by Z at $X = 0$, h represents the test of moderation of the M to Y path by X at $Z = 0$, and j represents the possibility that X and Z combine to moderate the M to Y path, an effect not considered by Edwards and Lambert or Preacher et al. Note also that c'_3 represents the moderated direct effect emphasized by Edwards and Lambert. Although MacKinnon (2008) presents the most general framework, he does not discuss each of the possible submodels in the depth of Edwards and Lambert or Preacher et al. MacKinnon also makes a clear distinction between the individual tests of the X to M and M to Y paths involved in the indirect effect and the tests of the overall indirect effect: They test different hypotheses.

In the present study we draw on these three general approaches to test a hypothesized model in which the direct relationship between symptoms and quality of life and indirect relationship between these two variables through disability may be moderated by age. We also utilize graphical and statistical procedures extended from Aiken and West (1991) to provide a further understanding of the effects we obtain.

Hypotheses

Mediation—The hypothesized basic (i.e., unconditional) mediation model is depicted in Figure 2. We hypothesized that the effect of Symptoms on Quality of life (QOL) would be partially mediated by Functional health (Disability). We expected that there would be an indirect relationship in which the changes in the level of Symptoms would be related to the level of Disability, which, in turn, would be related to QOL. We also expected that Symptoms would have a direct effect on QOL in addition to its effect through Disability. We expected that this general pattern of direct and indirect relationships would exist over the full range of ages represented in our adult sample.

Moderated direct and indirect effects—In addition to the unconditional mediation model, we hypothesized that the direct relationship of Symptoms to QOL and the relationship of Disability to QOL would be moderated by age, such that the relationships would be *stronger* for older relative to younger patients. The moderated direct and indirect effects model is shown in Figure 3. A large number of studies have shown relationships between age and symptom severity in various populations with medical conditions (e.g., cancer, Walsh, Donnelly, & Rybicki, 2000; depression, Prince et al., 1999; premenstrual symptoms, Freeman, Rickels, Schweizer, & Ting, 1995). A smaller number of studies have investigated relationships between age and quality of life, including one showing a negative relationship in HIV patients (Piette, Wachtel, Mor, & Mayer, 1995). However, with one exception, we were unable to locate any studies that have investigated age as a putative moderator of the *relationships* between symptoms, functional health, and quality of life. For the relationship of Disability to QOL, Masoudi et al. (2004) found in a population of heart disease patients that decreased functional health was associated with declines in health-related QOL in the older (age > 65) but not in the younger patients.

Methods

Data and measures

The data were collected from HIV positive patients recruited through clinics in San Francisco, Los Angeles, and San Diego. The sample we used in this analysis was $N = 956$ males collected between 1992 and 1994 prior to the introduction of the current treatment

regimen for HIV introduced in 1998. All participants had complete responses on Symptoms, Functional Health, and Health-Related Quality of Life items. Symptoms were measured using a 23 item checklist indicating whether each symptom occurred during the past week (e.g., have had fever, night sweats, nausea) with the response options being “yes” or “no.” Functional Health (Disability) was measured by 8 items asking whether they have had any difficulty carrying out daily tasks or activities (e.g., dress yourself, open a new carton of milk, climb up five steps). Disability items were scaled 0 (without any difficulty) to 3 (unable to do) so that higher numbers indicated more difficulty. Health-Related Quality of Life (QOL) was measured by 4 subscales based on 17 items measuring cognition (e.g., react slowly), vitality (e.g., feel worn out), mental health (e.g., feel nervous), and disease worry (e.g., frustrated about his health). QOL subscales were scaled from 1 to 5 with higher numbers indicating better quality of life.

We took the average of the 23 symptom items and the 8 functional health items to compute composite scores for Symptoms and Disability, respectively. We took the average of the 4 QOL subscale scores to compute composite scores for QOL. Coefficient alpha (Cronbach, 1951) was .876, .913, and .937 for Symptoms, Disability, and QOL, respectively. Previous work by Sousa and colleagues on this sample of patients has established the measurement structure underlying each of the three scales. Sousa, Tann, and Kwok (2006) showed that a second order factor model with a strong general second order factor (and six first order factors) fit the data well for the measure of symptoms in the HIV population. Sousa, Kwok, and Ryu (2008) showed that a single general factor underlies the functional health items in the HIV population. Chen, West, and Sousa (2006) showed a bifactor model with a strong general factor and much smaller specific factors associated with each subscale fit these data well in the HIV population. The means, standard deviations, and correlations of Symptoms, Disability, QOL, and Age are shown in Table 1. Following Aiken and West (1991), all of the composite scores for the independent variables (Symptoms, Disability, and Age) were centered at the mean. QOL was not centered.

Models

Mediation—The hypothesized mediational model presented in Figure 2 is expressed by the two regression equations as shown in (7) and (8). Recall that we expected that direct and indirect relationships would exist over the full range of ages represented in our adult sample.

$$M = a_{11} + b_{1M,X}X + e_{11} \quad (7)$$

$$Y = a_{12} + b_{1Y,X}X + b_{1Y,M}M + e_{12} \quad (8)$$

Substituting (7) into (8) yields the reduced form equation (9).

$$Y = (a_{12} + b_{1Y,M}a_{11}) + (b_{1Y,X} + b_{1Y,M}b_{1M,X})X + b_{1Y,M}e_{11} + e_{12} \quad (9)$$

In equation (7), the effects of X (Symptoms) on Y (QOL) consist of two additive components: the direct effect ($b_{1Y,X}$) and the indirect effect ($b_{1Y,M}b_{1M,X}$). The indirect effect is the product of the path coefficient from X (Symptoms) to M (Disability) and the path coefficient from M (Disability) to Y (QOL). We used three different methods to test the significance of the indirect effect: Sobel (1982) standard error, bias-corrected bootstrap confidence interval

(MacKinnon, Lockwood, & Williams, 2004), and asymmetric confidence interval produced by PRODCLIN program (MacKinnon, Fritz, Williams, & Lockwood, 2007). The three methods are reported for the purpose of illustration. In practice, researchers should choose one of the methods depending on the availability of the methods for their mediational model. The bias-corrected bootstrap method and the PRODCLIN program produce asymmetric confidence limits of the indirect effect. Both methods are more exact and have greater statistical power than those based on the normal distribution (e.g., Sobel method). For the two-path mediated effects, PRODCLIN may have trivially greater power than the bias-corrected bootstrap method. The bias-corrected bootstrap method also requires raw data which may sometimes be unavailable; it can also be computationally cumbersome. However, the PRODCLIN confidence limits cannot presently be computed for indirect effects consisting of more than two paths (e.g., three-path mediated effect: $X \rightarrow M1 \rightarrow M2 \rightarrow Y$), whereas the bias-corrected bootstrap method can be extended to more complicated cases. Although the least accurate, the Sobel method is easy to compute as it is now implemented in nearly all structural equation modeling software.

Moderated direct and indirect effects—The hypothesized moderated direct and indirect effects model presented in Figure 3 is expressed by the two regression equations (10) and (11). We tested the moderating effects of Age by adding the interaction terms ($XZ = \text{Symptoms} \times \text{Age}$ and $MZ = \text{Disability} \times \text{Age}$) as shown in (11).

$$M = a_{21} + b_{2M,X}X + e_{21} \quad (10)$$

$$Y = a_{22} + b_{2Y,X}X + b_{2Y,M}M + b_{2Y,Z}Z + b_{2Y,XZ}XZ + b_{2Y,MZ}MZ + e_{22} \quad (11)$$

Substituting (10) into (11) gives (12), in which the effects of X (Symptoms) on Y (QOL) consist of two additive components: the direct effect ($b_{2Y,X} + b_{2Y,XZ}Z$) and the indirect effect ($b_{2Y,M} + b_{2Y,MZ}Z$).

$$Y = [a_{22} + b_{2Y,M}a_{21} + (b_{2Y,Z} + b_{2Y,MZ}a_{21})Z] + [(b_{2Y,X} + b_{2Y,XZ}Z) + b_{2M,X}(b_{2Y,M} + b_{2Y,MZ}Z)]X + (b_{2Y,M} + b_{2Y,MZ}Z)e_{21} + e_{22} \quad (12)$$

The direct effect is a function of the moderating variable Age. If $b_{2Y,XZ}$ is significant, it means that the strength of the direct effect of Symptoms on QOL differs depending on how old the patient is. The indirect effect is also a function of the moderating variable Age. If $b_{2Y,MZ}$ is significant, it indicates that the Disability \rightarrow QOL relationship differs depending on the patient's age. The significance of the moderation of Age on the indirect effect (Symptoms \rightarrow Disability \rightarrow QOL) can be tested by testing the significance of the product term $b_{2M,X}b_{2Y,MZ}$. We obtained the bias-corrected bootstrap confidence interval to test the significance by utilizing the SAS macro provided by Taylor, MacKinnon, and Tein (2008). We used Mplus 5 to estimate the model.

Results

Unconditional Mediation Model—Using Sobel's (1982) standard error, both the direct effect (unstandardized estimate $\hat{b}_{1Y,X} = -2.028$, s.e. = 0.167, $p < .001$) and the indirect effect (unstandardized estimate $\hat{b}_{1Y,M}\hat{b}_{1M,X} = -0.217 \times 0.809 = -0.176$, s.e. = 0.048, $p < .001$) of Symptoms on QOL were significant. The estimates indicated that patients having higher scores on the symptoms checklist had lower scores on QOL. As noted above, the Sobel

method for testing the indirect effect is asymptotically correct, but fails to represent the asymmetric non-normal distribution of the product term at realistic sample sizes. In practice, this typically means that the Sobel method will have reduced power to test the indirect effect that may be appreciable with small effect sizes or small sample sizes (MacKinnon et al., 2002). Following recommendations by Edwards and Lambert and Preacher et al., we also used the bias-corrected bootstrapping method to construct the 95% confidence interval for the indirect effect. The 95% confidence interval for the indirect effect was $[-0.309, -0.039]$ and did not include zero. We can conclude that the indirect effect is significantly different from zero at $\alpha = .05$. Finally, we also used MacKinnon, Fritz, Williams, and Lockwood's (2007) PRODCLIN program which implements work by Aroian (1944; see also Meeker, Cornwell, & Aroian, 1981) on the distribution of the product of two normally distributed random variables. PRODCLIN produces asymmetric confidence intervals for the indirect effect at a specified level of $1 - \alpha$. If the confidence interval does *not* contain zero, the null hypothesis that the indirect effect equals 0 can be rejected at the corresponding level of α . In the present case, PRODCLIN produced lower and upper 95% confidence limits of -0.301 and -0.061 ; we can again conclude there is a significant indirect effect at $\alpha = .05$. All three methods indicated that the indirect effect of Symptoms on QOL through Disability was significantly different from zero.

The negative sign of the estimate of the direct effect ($\hat{b}_{1Y,X} = -2.028$) indicates that patients who have more symptoms have lower QOL. The significant indirect effect ($\hat{b}_{1Y,M}\hat{b}_{1M,X} = -0.176$) indicates that in addition to the direct effect of Symptoms on QOL, there is a second, independent process in which Symptoms is related to increasing Disability ($\hat{b}_{1M,X} = 0.809$, positive), which, in turn, is related to lower QOL ($\hat{b}_{1Y,M} = -0.217$, negative). Approximately 8% of the total effect of Symptoms on QOL is mediated through Disability.

Moderated direct and indirect effects—Recall that we hypothesized that age would moderate two of the paths in our mediational model. As shown in Figure 3, Age significantly moderated the direct effect of Symptoms on QOL ($\hat{b}_{2Y,XZ} = -0.031$, *s.e.* = 0.005, $p < .001$). The estimate indicated that the direct negative relationship of Symptoms to QOL became stronger by 0.031 for each 1 year increase in Age. In contrast, contrary to our hypothesis, we found no evidence for a moderating effect of age on the relationship between Disability and QOL to reach statistical significance ($\hat{b}_{2Y,MZ} = -0.003$, *s.e.* = 0.007, *ns*). The estimated moderating effect of Age on the indirect effect was $-0.002 (= 0.809 \times (-0.003))$. The bias-corrected 95% confidence interval was $[-0.007, 0.021]$. Therefore the moderation of Age on the indirect effect was also not significant.

In order to understand how the direct effect of Symptoms on QOL varies at different Age values, we estimated simple (conditional) slopes for the direct effect of Symptoms on QOL at different levels of Age (sample range = 21 to 69). We chose values of Age of 24, 32, 40, 48, and 56, which represent approximate values of the Mean Age $-2SD$, $M - 1SD$, M , $M + 1SD$, and $M + 2SD$, respectively. Following procedures described by Aiken and West (1991) and by Preacher, Curran, and Bauer (2006), we tested each of the conditional slopes. As shown in Figure 4, the estimated simple slopes were negative and statistically significant at all Age values ($-1.590, -1.838, -2.086, -2.334, \text{ and } -2.582$ at age = 24, 32, 40, 48, and 56, respectively; all $ps < .05$); the simple slopes became increasingly negative as Age increased. Otherwise stated, higher levels of Symptoms were associated with lower QOL at *all* ages, but this relationship became stronger as age increased. Indeed, when we used the extension of the Johnson-Neyman procedure (Preacher, Curran, & Bauer, 2006) to calculate the range of ages over which the conditional slopes between symptoms and quality of life is statistically significant, we found that this relationship was significant at the full range of values of age included in the study. Solution of the Johnson-Neyman equation showed

mathematically that the effect became nonsignificant at age = -21, representing an impossible value of age.

Discussion

Symptom status was found to be negatively related to the quality of life (QOL) of HIV patients. This relationship is partially mediated by the functional health (Disability) in that having more symptoms increased the levels of Disability, which, in turn, lowered QOL. The direct relationship between Symptoms and QOL was also significant. We also tested whether the direct and indirect effects of Symptoms on QOL varied depending on Age (i.e., were moderated by Age). We found that only the direct effect was moderated by Age. The relationship between Symptoms and QOL was negative with a higher level of Symptoms being associated with a lower QOL at all ages. However, the magnitude of the negative relationship became stronger in older patients. One possible explanation is that the patients' coping strategies decrease and their social support diminishes as they age. Studies have shown that active coping and social support are associated with good quality of life in patients with HIV (Friedland, Renwick, & McColl, 1996; Leiberich et al., 1997; Swindells et al., 1999). However, studies also have shown that older persons with HIV have limited social support and coping strategies (Heckman et al., 2002; Pitts, Grierson, & Misson, 2005).

The moderated path analysis approach presented here based on work by Edwards and Lambert (2007), Preacher et al. (2008), and MacKinnon (2008) has advantages over traditional approaches to moderated mediation. We were able to develop focused tests of a priori hypotheses about the forms of moderation of the direct and indirect effects that might be present. This ability contrasts with traditional approaches to moderated mediation in three important ways.

First, the traditional moderated causal steps procedure (Baron & Kenny, 1986; Muller, Judd, & Yzerbyt, 2005) can provide only a binary decision whether the combined mediation-moderation effect is present. The general approach illustrated here provides estimates of the direct and indirect effects, and it shows how these effects vary as a function of the moderator. In our data, we did not find a moderated indirect effect, but bootstrapping methods described by Edwards and Lambert (2007) and Preacher et al. (2007) are available to provide tests of conditional indirect effects.

Second, some traditional models have combined mediation and moderation by adding the moderator (Z) and the product of the independent variable \times moderator (XZ), as if there are three independent variables in the model (Baron & Kenny, 1986; Morgan-Lopez & MacKinnon, 2006; Morgan-Lopez, Castro, Chassin, & MacKinnon, 2003). In this model, which has often been termed *mediated moderation* model, the three independent variables (X , Z , and XZ) predict both M and Y . However, this mediated moderation model is equivalent to a particular case of the moderated path analysis model in which the $X \rightarrow M$ path and the $X \rightarrow Y$ path are *both* moderated by Z . In contrast, the use of approach described in this manuscript has greater flexibility in that it permits tests of moderation in which the $X \rightarrow M$ path, $M \rightarrow Y$ path, or $X \rightarrow Y$ path or any combination thereof are hypothesized to be moderated by Z . In addition, both the Preacher et al. and MacKinnon approaches permit X to moderate the $M \rightarrow Y$ path.

Third, particularly in the Edwards and Lambert approach, the moderated direct effect, here an effect of central interest, is emphasized as well as the moderated indirect effect. Traditional approaches to combining mediation and moderation have disregarded the direct effect. Preacher et al. also do not consider this type of effect and MacKinnon et al. include it

in their general model, but do not develop its possibilities. When there is a partial mediation, the relationship between the two variables is accounted for by two additive components – direct and indirect effects. In our analysis, the moderated direct effect has an important clinical implication. The impact of symptoms on QOL is greater for older patients. Therefore, symptom management is more crucial for older patients to help them maintain good quality of life.

We caution that our findings are based on a cross-sectional analysis like many other mediational analyses. Maxwell and Cole (2007) note that estimates based on cross-sectional data may be biased relative to estimates based on longitudinal data sets because of the failure to allow for autoregressive effects of the mediator and outcome variables. Baron and Kenny (1986), Holland (1988), and MacKinnon (2008) all note that in the absence of a randomized experiment we cannot rule out the possibility that other alternative explanations may also account for the present results. However in the present context manipulation of chronic symptom status and disability status is clearly precluded by important ethical concerns. Fortunately, in the present case some prominent alternative models are implausible: for example, current theory and research does not provide an account suggesting how quality of life could produce disability or symptoms. Another issue is that the present results imply a comparison between younger and older cohorts of patients, not the change of the relationship between Symptoms and QOL within patients as they age. Further research using focused tests associated with the moderated path analysis approach on longitudinal data sets can help us better understand and reach stronger conclusions about some of the mechanisms involved in the relationship between symptom status and quality of life in HIV patients.

Acknowledgments

This research was supported by National Institute of Nursing Research Grant NRO4817 to Karen Sousa and National Institute on Drug Abuse Grant DA09757 to David MacKinnon.

References


- Aiken, LS.; West, SG. Multiple regression: Testing and interpreting interactions. Sage Publications, Inc; Thousand Oaks, CA: 1991.
- Alwin DF, Hauser RM. The decomposition of effects in path analysis. *American Sociological Review* 1975;40:35–47.
- Aroian LA. The probability function of the product of two normally distributed variables. *Annals of Mathematical Statistics* 1944;18:265–271.
- Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology* 1986;51:1173–1182. [PubMed: 3806354]
- Bollen, KA. Total, direct and indirect effects in structural equation models. In: Clogg, CC., editor. *Sociological Methodology* 1987. American Sociological Association; Washington, DC: 1987. p. 37-69.
- Chaplin, WF. Moderator and mediator models in personality research: A basic introduction. In: Robins, RW.; Fraley, RC.; Krueger, R., editors. *Handbook of research methods in personality psychology*. Guilford Press; New York: 2007. p. 602-632.
- Chen FF, West SG, Sousa KH. A comparison of bifactor and second-order models of quality of life. *Multivariate Behavioral Research* 2006;41:189–225.
- Cronbach LJ. Coefficient alpha and the internal structure of tests. *Psychometrika* 1951;16:297–334.
- Cunningham WE, Crystal S, Bozzette S, Hays RD. The association of health-related quality of life with survival among persons with HIV infection in the United States. *Journal of General Internal Medicine* 2005;20:21–27. [PubMed: 15693923]

- Deng S, Lopez V, Roosa MW, Ryu E, Burrell GL, Tein JY, Crowder S. Family processes mediating the relationship of neighborhood disadvantage to early adolescent internalizing problems. *Journal of Early Adolescence* 2006;26:206–231.
- Edwards JR, Lambert LS. Methods for integrating moderation and mediation: A general analytical framework using moderated path analysis. *Psychological Methods* 2007;12:1–22. [PubMed: 17402809]
- Fox J. Effect analysis in structural equation models. *Sociological Methods and Research* 1980;9:3–26.
- Freeman EW, Rickels K, Schweizer E, Ting T. Relationships between age and symptom severity among women seeking medical-treatment for premenstrual symptoms. *Psychological Medicine* 1995;25:309–315. [PubMed: 7675918]
- Friedland J, Renwick R, McColl MM. Coping and social support as determinants of quality of life in HIV/AIDS. *AIDS Care* 1996;8:15–31. [PubMed: 8664366]
- Fritz MS, MacKinnon DP. Required sample size to detect the mediated effect. *Psychological Science* 2007;18:233–239. [PubMed: 17444920]
- Heckman TG, Heckman BD, Kochman A, Sikkema KJ, Suhr J, Goodkin K. Psychological symptoms among persons 50 years of age and older living with HIV disease. *Aging & Mental Health* 2002;6:121–128. [PubMed: 12028880]
- Holland PW. Causal inference, path analysis, and recursive structural equation modeling. *Sociological Methodology* 1988;18:449–484.
- Johnson PO, Neyman J. Tests of certain linear hypotheses and their applications to some educational problems. *Statistical Research Memoirs* 1936;1:57–93.
- Judd CM, Kenny DA. Process analysis: Estimating mediation in treatment evaluations. *Evaluation Review* 1981;5:602–619.
- Kenny, DA.; Kashy, DA.; Bolger, N. Data analysis in social psychology. In: Gilbert, D.; Fiske, S.; Lindzey, G., editors. *The handbook of social psychology*. 4th ed.. Vol. 1. McGraw-Hill; Boston, MA: 1998. p. 233-265.
- Kraemer HC, Wilson GT, Fairburn CG, Agras WS. Mediators and moderators of treatment effects in randomized clinical trials. *Archives of General Psychiatry* 2002;59:877–883. [PubMed: 12365874]
- Leiberich P, Engeter M, Olbrich E, Rubbert A, Schumacher K, Brieger M, Kalden JR, Joraschky P. Longitudinal development of distress, coping and quality of life in HIV-positive persons. *Psychotherapy and Psychosomatics* 1997;66:237–247. [PubMed: 9311027]
- MacCallum RC, Zhang S, Preacher KJ, Rucker DD. On the practice of dichotomization of quantitative variables. *Psychological Methods* 2002;7:19–40. [PubMed: 11928888]
- MacKinnon, DP. *Introduction to statistical mediation analysis*. Erlbaum; Mahwah, NJ: 2008.
- MacKinnon DP, Fritz MS, Williams J, Lockwood CM. Distribution of the product confidence limits for the indirect effect: Program PRODCLIN. *Behavior Research Methods* 2007;39:384–389. [PubMed: 17958149]
- MacKinnon DP, Lockwood CM, Hoffman JM, West SG, Sheets V. A comparison of methods to test mediated and other intervening variable effects. *Psychological Methods* 2002;7:83–104. [PubMed: 11928892]
- MacKinnon DP, Lockwood CM, Williams J. Confidence limits for the indirect effect: Distribution of the product and resampling methods. *Multivariate Behavioral Research* 2004;30:41–62. [PubMed: 20157641]
- MacKinnon DP, Warsi G, Dwyer JH. A simulation study of mediated effect measures. *Multivariate Behavioral Research* 1995;30:41–62. [PubMed: 20157641]
- Masoudi FA, Rumsfeld JS, Havranek EP, House JA, Peterson ED, Krumholz HM, Spertus JA. Age, functional capacity, and health-related quality of life in patients with heart failure. *Journal of Cardiac Failure* 2004;10:368–373. [PubMed: 15470645]
- Maxwell SE, Cole DA. Bias in cross-sectional analyses of longitudinal mediation. *Psychological Methods* 2007;12:23–44. [PubMed: 17402810]
- McBride C, Atkinson L, Quilty LC, Bagby RM. Attachment as moderator of treatment outcome in major depression: A randomized control trial of interpersonal psychotherapy versus cognitive behavior therapy. *Journal of Consulting and Clinical Psychology* 2006;74:1041–1054. [PubMed: 17154734]

- McNemar Q. At random: Sense and nonsense in psychology. *American Psychologist* 1960;15:295–300.
- Meeker, WQ., Jr.; Cornwell, LW.; Aroian, LA. The product of two normally distributed random variables. In: Kennedy, WJ.; Odeh, RE., editors. *Selected tables in mathematical statistics, Volume VII*. American Mathematical Society; Providence, RI: 1981.
- Morgan-Lopez AA, Castro FG, Chassin L, MacKinnon DP. A mediated moderation model of cigarette use among Mexican American youth. *Addictive Behaviors* 2003;28:583–589. [PubMed: 12628629]
- Morgan-Lopez AA, MacKinnon DP. Demonstration and evaluation of a method assessing mediated moderation. *Behavior Research Methods* 2006;38:77–87. [PubMed: 16817516]
- Muller D, Judd CM, Yzerbyt VY. When moderation is mediated and mediation is moderated. *Journal of Personality and Social Psychology* 2005;89:852–863. [PubMed: 16393020]
- Ostrander R, Crystal DS, August G. Attention deficit-hyperactivity disorder, depression, and self- and other-assessments of social competence: A developmental study. *Journal of Abnormal Child Psychology* 2006;34:773–787. [PubMed: 17066222]
- Piette J, Wachtel TJ, Mor V, Mayer K. The impact of age on the quality of life in persons with HIV infection. *Journal of Aging and Health* 1995;7:163–178. [PubMed: 10165953]
- Pitts M, Grierson J, Misson S. Growing older with HIV: A study of health, social and economic circumstances for people living with HIV in Australia over the age of 50 years. *AIDS Patient Care and STDs* 2005;19:460–465. [PubMed: 16053403]
- Preacher KJ, Curran PJ, Bauer DJ. Computational tools for probing interaction effects in multiple linear regression, multilevel modeling, and latent curve analysis. *Journal of Educational and Behavioral Statistics* 2006;32:437–448.
- Preacher KJ, Rucker DD, Hayes AF. Addressing moderated mediation hypotheses: Theory, methods, and prescriptions. *Multivariate Behavioral Research* 2007;42:185–227.
- Prince MJ, Beekman ATF, Deeg DJH, Fuhrer R, Kivela SL, Lawlor BA, Lobo A, Magnusson H, Meller I, Van Oyen H, Reischies F, Roelands M, Skoog I, Turrina C, Copeland JRM. Depression symptoms in late life assessed using the EURO-D scale: Effect of age, gender and marital status in 14 European centers. *British Journal of Psychiatry* 1999;174:339–345. [PubMed: 10533553]
- Roosa MW, Deng S, Ryu E, Burrell G,L, Tein JY, Jones S, Lopez V, Crowder S. Family and children characteristics linking neighborhood context and child externalizing behavior. *Journal of Marriage and Family* 2005;67:515–529.
- Sobel, ME. Asymptotic confidence intervals for indirect effects in structural equation models. In: Leinhardt, S., editor. *Sociological methodology*. Jossey-Bass; San Francisco: 1982. p. 290-312.
- Shrout PE, Bolger N. Mediation in experimental and , nonexperimental studies: New procedures and recommendations. *Psychological Methods* 2002;7:422–445. [PubMed: 12530702]
- Sousa KH, Holzemer WL, Henry SB, Slaughter R. Dimensions of health-related quality of life in persons living with HIV disease. *Journal of Advanced Nursing* 1999;29:178–187. [PubMed: 10064297]
- Sousa KH, Kwok O. Examining the longitudinal impact of symptom status on QOL: A parallel process growth modeling [Abstract]. *Quality of Life Research* 2005;14:2018.
- Sousa KH, Kwok O. Putting Wilson and Cleary to the test: Analysis of a HRQOL conceptual model using structural equation modeling. *Quality of Life Research* 2006;15:725–737. [PubMed: 16688505]
- Sousa KH, Kwok O, Ryu E. Confirmation of the validity of the HAQ-DI in two populations living with chronic illnesses. *Journal of Nursing Measurement* 2008;16:31–42. [PubMed: 18578108]
- Sousa KH, Tann SS, Kwok O. Reconsidering the assessment of symptoms status in HIV/AIDS care. *Journal of the Association of Nurses in AIDS Care* 2006;17:36–46. [PubMed: 16800166]
- Swindells S, Mohr J, Justis JC, Berman S, Squier C, Wagener MM, Singh N. Quality of life in patients with human immunodeficiency virus infection: Impact of social support, coping style, and hopelessness. *International Journal of STD & AIDS* 1999;10:383–391. [PubMed: 10414881]
- Taylor AB, MacKinnon DP, Tein JY. Test of the three-path mediated effect. *Organizational Research Methods* 2008;11:241–269.


- Walsh D, Donnelly S, Rybicki L. The symptoms of advanced cancer: relationship to age, gender, and performance status in 1000 patients. *Supportive Care in Cancer* 2000;8:175–179. [PubMed: 10789956]
- Wilson IB, Cleary PD. Linking clinical variables with health-related quality of life: A conceptual model of patient outcomes. *Journal of the American Medical Association* 1995;273:59–65. [PubMed: 7996652]
- Wilson IB, Cleary PD. Clinical predictors of declines in physical functioning in persons with AIDS: Results of a longitudinal study. *Journal of Acquired Deficiency Syndromes* 1997;16:343–349.

(A)



$$Y = i_1 + cX + e_1$$

(B)



$$M = i_2 + aX + e_2$$

$$Y = i_3 + c'X + bM + e_3$$

Figure 1.
Mediation model (Baron & Kenny, 1986; Judd & Kenny, 1981)




Figure 2. Mediation: Effects of Symptoms on QOL partially mediated by Disability. Unstandardized estimates are shown. * indicates $p < .05$.




Figure 3. Moderated direct and indirect effects: Direct effect of Symptoms on QOL moderated by Age. Unstandardized estimates are shown. * $p < .05$. ns = non-significant. Dashed arrow represents non-significant paths.




Figure 4. Estimated direct effects of Symptoms on QOL at different Age values. Each of the conditional direct effects is significant at $p < .05$.

Table 1

Means, standard deviations, and correlations of Symptoms, Disability, QOL, and Age

	Mean	SD	Correlation			
			Symptoms	Disability	QOL	Age
Symptoms	0.208	0.208	1.000			
Disability	0.147	0.275	0.611	1.000		
QOL	3.306	0.683	-0.669	-0.464	1.000	
Age	39.386	8.141	-0.113	-0.032	0.113	1.000

Note. N = 956.