

# Inflammatory Markers and Risk of Developing Type 2 Diabetes in Women

Frank B. Hu,<sup>1,2,3</sup> James B. Meigs,<sup>4</sup> Tricia Y. Li,<sup>1</sup> Nader Rifai,<sup>5</sup> and JoAnn E. Manson<sup>2,3,6</sup>

**We conducted a prospective, nested, case-control study of inflammatory markers as predictors of type 2 diabetes among 32,826 women who provided blood samples in 1989 through 1990 in the Nurses' Health Study. Among women free of diabetes, cardiovascular disease, or cancer at baseline, 737 had developed diabetes by 2000. Control women ( $n = 785$ ) were selected matched on age, fasting status, race, and BMI for cases in the top BMI decile. Baseline levels of tumor necrosis factor (TNF)- $\alpha$  receptor 2, interleukin (IL)-6, and C-reactive protein (CRP) were significantly higher among case than control subjects (all  $P \leq 0.001$ ). After adjusting for BMI and other lifestyle factors, all three biomarkers significantly predicted diabetes risk; the odds ratios (ORs) comparing extreme quintiles were 1.64 (95% CI 1.10–2.45) for TNF- $\alpha$ R2, 1.91 (1.27–2.86) for IL-6, and 4.36 (2.80–6.80) for CRP ( $P$  for trend  $<0.001$  for all biomarkers). In a multivariate model simultaneously including the three biomarkers, only CRP levels were significantly associated with risk of diabetes (OR comparing extreme quintiles of CRP = 3.99,  $P$  for trend  $<0.001$ ). These data support the role of inflammation in the pathogenesis of type 2 diabetes. Elevated CRP levels are a strong independent predictor of type 2 diabetes and may mediate associations of TNF- $\alpha$ R2 and IL-6 with type 2 diabetes. *Diabetes* 53:693–700, 2004**

**E**xcess adiposity is the most important risk factor for the development of insulin resistance and type 2 diabetes (1). However, mechanisms whereby body fat induces insulin resistance in distant tissues are not well understood. Recent evidence indicates that obesity may be an inflammatory condition (2). It has been proposed that inflammatory cytokines secreted by adipose tissue exert an endocrine effect con-

ferring insulin resistance in liver, skeletal muscle, and vascular endothelial tissue, ultimately leading to the clinical expression of both type 2 diabetes and cardiovascular disease (CVD) (3,4). In particular, elevated production of adipocyte cytokines, such as tumor necrosis factor (TNF)- $\alpha$  and interleukin (IL)-6, leads to an acute-phase response with increased hepatic production of C-reactive protein (CRP), a sensitive marker of low-grade systemic inflammation (5–7). TNF- $\alpha$ , IL-6, and CRP not only directly promote insulin resistance, but also stimulate endothelial production of adhesion molecules such as E-selectin, intercellular adhesion molecule-1 (ICAM-1), and vascular adhesion molecule-1 (VCAM-1), critical mediators of endothelial dysfunction in capillary and arteriolar endothelium (8).

Most previous epidemiological research has focused on the role of CRP in predicting risk of type 2 diabetes. While several studies observed a significant positive association between CRP levels and type 2 diabetes after adjusting for BMI or waist circumference (9–12), others suggested that the association was largely explained by obesity (13–16). In this large prospective, nested, case-control study, we simultaneously examined the role of TNF- $\alpha$  receptor 2 (TNF- $\alpha$ R2, as a measure of TNF- $\alpha$  in plasma), IL-6, and CRP in predicting the development of type 2 diabetes independent of obesity, diet, and other lifestyle factors in women.

## RESEARCH DESIGN AND METHODS

**Study population.** The Nurses' Health Study cohort was established in 1976 when 121,700 female registered nurses aged 30–55 years and residing in 11 U.S. states responded to mailed questionnaires regarding their medical history and health practices. Since 1976, questionnaires have been administered biennially to update health information and to identify new cases of disease. During 1989 through 1990, 32,826 women free of diagnosed diabetes, coronary heart disease, stroke, or cancer provided blood samples. By 2000, 737 of these women had developed definite diabetes. Control women providing baseline blood samples were matched to diabetes cases by year of birth, date of blood draw, race, and fasting status (at least 8 h overnight) at blood draw. In addition, to improve statistical control for obesity at the upper extreme of the distribution, control subjects were also matched on BMI with case subjects in the top 10% of the BMI distribution, giving a sample of 785 control women (26 control subjects were selected twice in the nested case-control risk-set sampling). The study was approved by the Institutional Review Board of the Brigham and Women's Hospital, Boston.

**Ascertainment of diabetes.** A validated supplementary questionnaire regarding symptoms, diagnostic tests, and hypoglycemic therapy was mailed to women who indicated on any biennial questionnaire that they had been diagnosed with diabetes. A case of diabetes was considered confirmed if at least one of the following was reported on the supplementary questionnaire: 1) classic symptoms plus elevated glucose levels (a fasting plasma glucose concentration  $\geq 140$  mg/dl or randomly measured concentration of at least 200 mg/dl); 2) at least two elevated plasma glucose concentrations on different occasions in the absence of symptoms (levels  $\geq 200$  mg/dl after 2 or more hours on oral glucose tolerance testing); 3) treatment with oral hypoglycemic

From the <sup>1</sup>Department of Nutrition, Harvard School of Public Health, Boston, Massachusetts; the <sup>2</sup>Channing Laboratory, Harvard Medical School and Brigham and Women's Hospital, Boston, Massachusetts; the <sup>3</sup>Department of Epidemiology, Harvard School of Public Health, Boston, Massachusetts; the <sup>4</sup>Department of Medicine, General Medicine Division, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts; the <sup>5</sup>Department of Pathology, Children's Hospital Medical Center and Harvard Medical School, Boston, Massachusetts; and the <sup>6</sup>Division of Preventive Medicine, Department of Medicine, Harvard Medical School and Brigham and Women's Hospital, Boston, Massachusetts.

Address correspondence and reprint requests to Dr. Frank Hu, Department of Nutrition, Harvard School of Public Health, 665 Huntington Ave., Boston, MA 02115. E-mail: frank.hu@channing.harvard.edu.

Received for publication 25 August 2003 and accepted in revised form 10 December 2003.

CRP, C-reactive protein; CVD, cardiovascular disease; ELISA, enzyme-linked immunosorbent assay; ICAM-1, intercellular adhesion molecule-1; IL, interleukin; TNF, tumor necrosis factor; TNF- $\alpha$ R2, TNF- $\alpha$  receptor 2; VCAM-1, vascular adhesion molecule-1.

© 2004 by the American Diabetes Association.

agents or insulin. These criteria are consistent with those proposed by the National Diabetes Data Group (NDDG) (17). The diagnostic criteria for type 2 diabetes were changed in 1997 (18) such that lower fasting glucose levels ( $>126$  mg/dl) would now be considered diagnostic. Thus, we have used the American Diabetes Association criteria for diagnosis of diabetes cases after 1998.

The validity of this diagnostic procedure has been verified in a subsample of this study population (19). The diagnosis of type 2 diabetes was confirmed by medical records in 98.4% (60 of 61) of the women. In addition, another substudy assessing the prevalence of undiagnosed diabetes suggested a very low rate of false negatives (20).

**Assessment of lifestyle factors.** Every 2 years, participant exposure status has been updated by questionnaire, including smoking (never smoked, former smoker, or current smoker), menopausal status, use or nonuse of postmenopausal hormone therapy, and body weight. Reported weights have been shown to correlate with measured weights ( $r = 0.96$ ) (21). We calculated BMI as weight in kilograms divided by the square of height (assessed in 1976) in meters. In 1986 we also assessed self-reported waist girth; this measure also strongly correlates with measured waist circumference ( $r = 0.89$ ) in this cohort (22). The presence or absence of a family history of diabetes in first-degree relatives was assessed in 1982 and 1988. Information about physical activity was assessed in 1988 using a validated questionnaire (23).

Diet was assessed using a validated semiquantitative food frequency questionnaire (24). A composite score was constructed on the basis of a diet low in *trans* fat and glycemic load, high in cereal fiber, and with a high ratio of polyunsaturated to saturated fat (25). For each dietary factor, we assigned each woman a score of one to five corresponding to her quintile of intake, with five representing the most favorable quintile, and summed her quintile values for the four nutrients.

**Laboratory procedures.** Women willing to provide blood specimens were sent instructions and a phlebotomy kit (including sodium heparin blood tubes, needles, and a tourniquet). Blood specimens were returned by overnight mail in a frozen water bottle, and upon arrival they were centrifuged to separate plasma from buffy coat and red cells and frozen in liquid nitrogen until analysis. Ninety-seven percent of samples arrived within 26 h of phlebotomy. Quality control samples were routinely frozen along with study samples to monitor for plasma changes due to long-term storage and to monitor assay variability. Previous work has documented the long-term stability of plasma samples collected and stored under this protocol (26). Study samples were analyzed in randomly ordered case-control pairs to further reduce systematic bias and interassay variation.

Frozen plasma aliquots from case and control subjects were selected for simultaneous analysis. CRP levels were measured via a high-sensitivity latex-enhanced immunonephelometric assay on a BN II analyzer (Dade Behring, Newark, DE). IL-6 was measured by a quantitative sandwich enzyme immunoassay technique (Quantikine HS Immunoassay kit) and TNF-R2 levels by an enzyme-linked immunosorbent assay (ELISA) kit utilizing immobilized monoclonal antibody to human TNF-R2 (Genzyme, Cambridge, MA). Levels of E-selectin and ICAM-1 were measured by commercial ELISA (R & D Systems, Minneapolis, MN). Insulin levels were measured with a double antibody system with  $<0.2\%$  cross-reactivity between insulin and its precursors (Linco Research, St. Louis, MO). HbA<sub>1c</sub> was measured by immunoassay (Hitachi 911 Analyzer; Roche Diagnostics, Indianapolis, IN). The coefficients of variation for each analyte were: 3.8% for C-reactive protein, 9.5% for fasting insulin, 3.8% for HbA<sub>1c</sub>, 5.9% for IL-6, 6.2% for TNF- $\alpha$ R2, and 6.6% for E-selectin.

**Statistical analysis.** Baseline characteristics of case and control subjects were compared by Student's *t* tests, Wilcoxon's rank-sum tests (for differences in median values), or  $\chi^2$  tests. Pearson correlation coefficients were used to evaluate associations among various biomarkers. We used conditional logistic regression to estimate the OR and 95% CIs for levels of biomarkers predicting type 2 diabetes, conditioning on matching factors such as age, fasting status, and race. We divided the distributions of the inflammatory markers in the control subjects into quintiles and used regression models to estimate the significance of trend in ORs across increasing biomarker quintiles and to estimate the OR of diabetes in each quintile using the lowest quintile as the referent category. In multivariate analyses, we adjusted for BMI (modeled as a continuously distributed covariate), further adjusted for family history of diabetes, smoking, alcohol intake, physical activity, postmenopausal hormone use (all modeled as categorical covariates), and the diet score. We then included CRP, IL-6, and TNF- $\alpha$ R2 simultaneously in the multivariate model to mutually adjust for each other. In a secondary analysis among women providing fasting blood samples, we also adjusted models for levels of fasting insulin and HbA<sub>1c</sub>. In addition, we conducted analyses of inflammatory markers and diabetes risk stratified on levels of diabetes risk factors including obesity, physical activity, and the diet score. Likelihood ratio tests were conducted to test statistical interactions between inflammatory biomarkers

and obesity, physical activity, regular aspirin use (approximately once per week or more), family history of diabetes, and diet score by comparing the  $-2$  log(likelihood) between two nested models, one with only the main effects and the other with both the main effects and interaction terms. SAS Statistical (27) was used in all of the analyses.

## RESULTS

As expected, women who developed diabetes during follow-up had significantly higher levels of BMI, waist circumference, and waist-to-hip ratio; had lower levels of physical activity and intake of cereal fiber; and were more likely to have a family history of diabetes than control subjects (Table 1). Case subjects were slightly more likely to use aspirin than control subjects. Among case subjects, levels of TNF- $\alpha$ R2, CRP, IL-6, ICAM-1, E-selectin, and fasting insulin were all significantly higher at baseline in women who developed diabetes than in those who remained nondiabetic during follow-up.

Among the healthy control subjects, BMI and waist circumference were significantly correlated with CRP levels, and the correlations between obesity and TNF- $\alpha$ R2, IL-6, and endothelial markers were somewhat weaker (Table 2). CRP was moderately but significantly correlated with TNF- $\alpha$ R2, IL-6, ICAM-1, and E-selectin.

Elevated levels of TNF- $\alpha$ R2, IL-6, and CRP measured at baseline were significantly associated with risk of incident type 2 diabetes. In crude analyses conditioned on matching covariates, the OR of diabetes comparing extreme quintiles were 2.59 for TNF- $\alpha$ R2, 3.25 for IL-6, and 7.08 for CRP ( $P$  for trend  $<0.001$  for all of the biomarkers, Table 3). These ORs were substantially attenuated but remained statistically significant after adjustment for BMI. Further adjustment for other diabetes risk factors, including family history of diabetes, smoking, alcohol intake, physical activity, diet, and hormone use, did not appreciably alter the ORs (Table 3). A recent study suggested a significant association between inflammation and diabetes among nonsmokers but not smokers (28). However, we did not observe such an interaction between smoking and CRP levels ( $P = 0.43$ ).

When TNF- $\alpha$ R2, CRP, and IL-6 were included simultaneously in the model, only CRP remained significant (ORs across quintiles were 1.0, 1.29, 1.59, 3.48, and 3.99,  $P$  for trend  $<0.001$ ). The ORs for TNF- $\alpha$ R2 and IL-6 were substantially attenuated and did not reach statistical significance (ORs comparing extreme quintiles were 1.25, 95% CI 0.82–1.91 for TNF- $\alpha$ R2 and 1.16, 0.75–1.79 for IL-6). Because these biomarkers are in the same causal pathway of systemic inflammation and diabetes, these results suggest that the effects of TNF- $\alpha$ R2 and IL-6 may be mediated through elevated CRP. When the effects of the inflammatory markers were evaluated jointly, the increased diabetes risk tended to be additive (all  $P$  values for multiplicative interactions  $>0.05$  between the biomarkers), with the highest risk of diabetes among women with the highest levels of CRP in combination with the highest levels of IL-6 (Fig. 1A) and TNF- $\alpha$ R2 (Fig. 1B). However, the stratified analyses indicate that the effects of CRP were more evident than those of IL-6 or TNF- $\alpha$ R2. In joint analyses, the effects of TNF- $\alpha$ R2 and IL-6 tended to be independent and additive (Fig. 1C).

In a previous analysis (29), we found that markers of endothelial dysfunction, especially E-selectin, significantly

TABLE 1  
Comparison of diabetes risk factors between case and control subjects at baseline\*

	Case	Control	P
<i>n</i>	737	785	
Age (years)	56.3 ± 6.9	56.2 ± 6.9	0.69
BMI (kg/m <sup>2</sup> )	30.3 ± 5.6	26.2 ± 6.1	<0.001
Physical activity (METs/week) <sup>†</sup>	12.3 ± 15.2	15.6 ± 27.2	0.004
Alcohol consumption (g/day)	3.7 ± 7.1	6.5 ± 9.0	<0.001
Waist circumference (in)	35.3 ± 4.9	31.3 ± 4.9	<0.001
Diet score <sup>‡</sup>	7.8 ± 2.7	8.2 ± 2.7	<0.001
Current smoking (%)	14.1	13.3	0.62
Postmenopausal status (%)	36.0	42.3	0.01
Current postmenopausal hormone use (%)	39.8	36.0	0.35
Family history of diabetes in first-degree relative (%)	46.8	21.0	<0.001
Race (Caucasian) (%)	94.8	94.3	0.68
Aspirin use ≥1/week	25.1	20.9	0.05
Biomarkers (median values)			
TNF-αR2 (pg/ml)	2,646.5	2,383.8	<0.001
CRP (mg/dl)	0.36	0.16	<0.001
IL-6 (pg/ml)	2.38	1.84	<0.001
E-selectin (pg/ml)	61.25	45.37	<0.001
Fasting insulin (uU/ml) <sup>§</sup>	11.96	8.27	<0.001

Data are means ± SE unless otherwise indicated. \*Case and control subjects were matched on age, fasting status, and race. <sup>†</sup>MET, metabolic equivalent; 1 MET hour is equivalent to energy expended by sitting quietly for 1 h. <sup>‡</sup>Intakes for *trans* fat, cereal fiber, glycemic load, and P-to-S (polyunsaturated fat-to-saturated fat) ratio were categorized into quintiles and for each participant, the quintile value for each nutrient (a higher quintile score represents a lower risk) was summed and the sum was recategorized into quintiles. <sup>§</sup>A total of 432 case and 398 control subjects had fasting insulin measurement.

predicted risk of type 2 diabetes. To examine whether the association between CRP and risk of diabetes was mediated through markers of endothelial dysfunction, we simultaneously included CRP and E-selectin in the multivariate model. The association for CRP was independent of levels of E-selectin (OR for the highest versus lowest quintile = 3.52, *P* for trend <0.001), and the elevated risk with increasing CRP and E-selectin levels appeared to be additive and independent of each other (Fig. 1D) (*P* for interaction = 0.55).

In stratified analyses, levels of TNF-αR2 (without adjustment for CRP and IL-6) significantly predicted diabetes risk among nonobese women, women with lower physical activity levels, women with a lower diet score, and women with no family history of diabetes (Table 4). However, only the test for interaction between obesity and TNF-αR2 was statistically significant (*P* = 0.01). The association between IL-6 and diabetes risk appeared to be stronger for women with BMI <30 kg/m<sup>2</sup> and those with lower physical activity levels, although the tests for interactions were not statistically significant. The association between CRP levels and risk of diabetes was significant in stratified analy-

ses according to levels of obesity, physical activity, the diet score, and family history of diabetes (Table 4).

In additional analyses, we adjusted for regular use (once a week or more) of aspirin and other nonsteroidal anti-inflammatory drugs and obtained similar results for CRP (OR comparing extreme quintiles = 4.33, 95% CI 2.78–6.74, *P* for trend <0.001). However, in stratified analyses (Fig. 2), the association between CRP levels was substantially stronger among participants who did not use aspirin (multivariate ORs across quintiles of CRP were 1.0, 1.13, 2.06, 4.86, and 8.85, *P* for trend <0.001) compared with women who reported regular aspirin use (corresponding ORs were 1.0, 1.49, 1.12, 3.33, and 2.94, *P* for trend <0.001, *P* for interaction between aspirin use and CRP = 0.0002).

We conducted several sensitivity analyses to address the issues of undiagnosed diabetes in the control subjects. First, we removed women with HbA<sub>1c</sub> levels >6.0% and repeated the analyses; the results remained virtually unchanged. Second, adjustment for HbA<sub>1c</sub> levels for both case and control subjects in the multivariate analyses had virtually no impact on the results (multivariate OR comparing extreme quintiles of CRP was 4.33, 95% CI 1.98–

TABLE 2  
Spearman correlation coefficients\* between BMI, waist circumference, fasting insulin, and inflammatory markers in control subjects

	BMI	Waist circumference	Fasting insulin	CRP	IL-6	TNF-αR2	E-selectin
BMI	1.0	—	—	—	—	—	—
Waist circumference	0.78	1.0	—	—	—	—	—
Fasting insulin	0.29	0.27	1.0	—	—	—	—
CRP	0.44	0.37	0.26	1.0	—	—	—
IL-6	0.30	0.29	0.18	0.37	1.0	—	—
TNF-αR2	0.24	0.25	0.15	0.27	0.28	1.0	—
E-selectin	0.28	0.27	0.26	0.24	0.25	0.23	1.0

\*All correlation coefficients were statistically significant at *P* < 0.05.

TABLE 3

RR (95% CI) of type 2 diabetes according to quintiles of inflammatory markers estimated from conditional logistic regression models

	Q1	Q2	Q3	Q4	Q5	<i>P</i> for trend
<b>TNF<math>\alpha</math>R2 (pg/ml)</b>						
Range	≤1,927.7	1,827.8–2,224.4	2,224.5–2,530.0	2,530.1–3,023.5	≥3,023.6	
Median	1,705.6	2,085.5	2,385.9	2,737.5	3,448.4	
<i>n</i>	96/167	98/155	124/153	192/155	227/155	
Crude RR*	1.00	1.14 (0.79, 1.65)	1.29 (0.90, 1.84)	2.10 (1.49, 2.95)	2.59 (1.84, 3.65)	<0.001
BMI-adjusted RR†	1.00	1.00 (0.68, 1.48)	1.11 (0.76, 1.62)	1.56 (1.08, 2.24)	1.67 (1.16, 2.42)	<0.001
Multivariate RR‡	1.00	0.89 (0.58, 1.37)	1.09 (0.73, 1.64)	1.47 (0.99, 2.18)	1.64 (1.10, 2.45)	0.001
<b>IL-6 (pg/ml)</b>						
Range	≤1.12	1.13–1.62	1.63–2.13	2.15–3.07	≥3.08	
Median	0.88	1.34	1.84	2.55	4.24	
<i>n</i>	88/185	110/150	134/150	176/151	229/149	
Crude RR	1.00	1.60 (1.11, 2.31)	1.77 (1.24, 2.53)	2.44 (1.71, 3.46)	3.25 (2.30, 4.58)	<0.001
BMI-adjusted RR	1.00	1.29 (0.87, 1.90)	1.21 (0.83, 1.78)	1.55 (1.06, 2.26)	1.95 (1.34, 2.84)	<0.001
Multivariate RR	1.00	1.24 (0.82, 1.88)	1.23 (0.81, 1.86)	1.43 (0.95, 2.15)	1.91 (1.27, 2.86)	0.001
<b>CRP (mg/dl)</b>						
Range	≤0.055	0.056–0.118	0.119–0.208	0.209–0.404	≥0.405	
Median	0.03	0.083	0.158	0.291	0.627	
<i>n</i>	55/176	58/152	89/153	216/152	319/152	
Crude RR	1.00	1.36 (0.87, 2.13)	2.11 (1.38, 3.22)	4.93 (3.33, 7.31)	7.08 (4.84, 10.37)	<0.001
BMI-adjusted RR	1.00	1.20 (0.75, 1.91)	1.68 (1.08, 2.62)	3.64 (2.42, 5.49)	4.18 (2.78, 6.29)	<0.001
Multivariate RR	1.00	1.30 (0.80, 2.13)	1.62 (1.01, 2.06)	3.61 (2.33, 5.60)	4.36 (2.80, 6.80)	<0.001

\*Conditioned on matching variables: age, fasting status, and race; †BMI as a continuous variable; ‡adjusted for time at blood drawn, alcohol consumption (nondrinkers, 0–4.9, 5–10, and >10 g/day), physical activity (quintiles of METs/week), smoking status (never, past, and current smoking of 1–14, 15–24, and ≥25 cigarettes per day), BMI (continuous), family history of diabetes, postmenopausal hormone use and menopausal status (never/unknown user or premenopausal status, past, and current user), and diet score (in quintiles).

9.44, *P* for trend <0.001). To ensure that the association of inflammatory biomarkers with incident diabetes was not due to subclinical CVD at baseline, we repeated the analysis after removing all women with incident CVD that developed during follow-up; again, results were un-

changed (multivariate OR comparing extreme quintiles of CRP 4.43, 2.83–6.93, *P* for trend <0.001).

To examine whether the associations between inflammatory markers and risk of diabetes were mediated through hyperinsulinemia, we conducted additional anal-




FIG. 1. A–D: Multivariate OR of type 2 diabetes according to joint classifications of plasma levels of inflammatory and endothelial markers. Adjusted for the same variables as in Table 3.

TABLE 4

Multivariate RR (95% CI) of type 2 diabetes according to quintiles of inflammatory markers stratified by BMI, physical activity, diet score, and family history of diabetes\*

	Q1	Q2	Q3	Q4	Q5	<i>P</i> for trend	<i>P</i> for interaction
<b>TNF<math>\alpha</math>R2</b>							
BMI <30 kg/m <sup>2</sup>	1.00	0.95 (0.58–1.57)	1.16 (0.72–1.87)	1.62 (1.02–2.59)	2.25 (1.39–3.62)	<0.001	
BMI $\geq$ 30 kg/m <sup>2</sup>	1.00	0.67 (0.26–1.68)	1.33 (0.53–3.35)	1.05 (0.45–2.44)	0.93 (0.41–2.09)	0.99	0.01
<b>Physical activity levels</b>							
Low	1.00	1.02 (0.58–1.77)	1.74 (1.02–2.98)	1.96 (1.18–3.28)	2.14 (1.29–3.54)	<0.001	
High	1.00	0.75 (0.41–1.38)	0.86 (0.48–1.54)	1.29 (0.74–2.26)	1.30 (0.72–2.34)	0.11	0.72
<b>Diet score</b>							
Low	1.00	0.74 (0.42–1.29)	1.04 (0.61–1.76)	1.45 (0.89–2.36)	2.05 (1.25–3.36)	<0.001	
High	1.00	1.06 (0.58–1.94)	1.32 (0.73–2.38)	1.78 (0.98–3.23)	1.18 (0.64–2.18)	0.48	0.21
<b>Family history of diabetes</b>							
Yes	1.00	0.73 (0.38–1.39)	0.75 (0.38–1.45)	1.15 (0.60–2.19)	1.51 (0.75–3.07)	0.07	
No	1.00	0.93 (0.54–1.62)	1.45 (0.87–2.41)	1.82 (1.12–2.96)	1.88 (1.16–3.04)	0.002	0.46
<b>IL-6</b>							
BMI <30 kg/m <sup>2</sup>	1.00	1.20 (0.75–1.92)	1.14 (0.70–1.84)	1.44 (0.90–2.30)	1.84 (1.16–2.94)	0.006	
BMI $\geq$ 30 kg/m <sup>2</sup>	1.00	0.90 (0.31–2.62)	1.11 (0.41–3.00)	1.07 (0.41–2.78)	1.46 (0.57–3.76)	0.17	0.29
<b>Physical activity levels</b>							
Low	1.00	1.31 (0.75–2.29)	1.27 (0.73–2.23)	1.70 (1.00–2.91)	2.47 (1.44–4.24)	<0.001	
High	1.00	1.38 (0.77–2.45)	1.63 (0.93–2.86)	1.44 (0.81–2.53)	1.77 (1.01–3.08)	0.09	0.73
<b>Diet score</b>							
Low	1.00	0.96 (0.56–1.65)	1.36 (0.80–2.29)	1.30 (0.78–2.16)	1.86 (1.12–3.09)	0.004	
High	1.00	1.74 (0.96–3.15)	1.28 (0.69–2.36)	1.85 (1.01–3.37)	2.16 (1.19–3.92)	0.024	0.38
<b>Family history of diabetes</b>							
Yes	1.00	1.99 (1.02–3.88)	1.79 (0.90–3.55)	2.19 (1.14–4.20)	1.97 (1.04–3.73)	0.16	
No	1.00	0.90 (0.54–1.50)	1.15 (0.70–1.87)	1.29 (0.80–2.10)	1.95 (1.21–3.13)	<0.001	0.13
<b>CRP</b>							
BMI <30 kg/m <sup>2</sup>	1.00	0.80 (0.46–1.37)	0.98 (0.57–1.69)	2.42 (1.49–3.95)	3.57 (2.13–5.96)	<0.001	
BMI $\geq$ 30 kg/m <sup>2</sup>	1.00	1.74 (0.48–6.27)	1.42 (0.49–4.18)	2.76 (1.01–7.58)	3.38 (1.31–8.77)	0.004	0.13
<b>Physical activity levels</b>							
Low	1.00	1.28 (0.69–2.37)	1.51 (0.83–2.74)	3.71 (2.14–6.43)	4.28 (2.47–7.43)	<0.001	
High	1.00	0.86 (0.42–1.78)	1.34 (0.68–2.62)	2.81 (1.49–5.30)	4.01 (2.09–7.68)	<0.001	0.87
<b>Diet score</b>							
Low	1.00	1.03 (0.55–1.92)	1.28 (0.72–2.27)	2.91 (1.70–5.01)	4.33 (2.50–7.49)	<0.001	
High	1.00	1.23 (0.61–2.49)	1.69 (0.83–3.46)	3.89 (2.05–7.36)	4.10 (2.13–7.88)	<0.001	0.94
<b>Family history of diabetes</b>							
Yes	1.00	1.49 (0.66–3.36)	1.84 (0.89–3.79)	3.27 (1.61–6.61)	3.73 (1.85–7.56)	<0.001	
No	1.00	0.94 (0.51–1.72)	1.15 (0.64–2.11)	3.30 (1.97–5.52)	4.66 (2.74–7.89)	<0.001	0.15

\*Physical activity and diet score were classified into low and high groups according to median values. RRs were adjusted for the same variables as in table 3.

yses adjusting for fasting insulin levels in the subset of women who provided fasting samples (432 case and 398 control subjects), and the association was only slightly attenuated (OR comparing extreme quintiles of CRP 3.85,

95% CI 2.08–7.13, *P* for trend <0.001). Finally, in the subset of women who provided waist circumference measures, we adjusted for both BMI and waist circumference (continuous variables); again, the association for CRP did not substantially change (OR comparing extreme quintiles of CRP 3.61, 2.29–5.68, *P* for trend <0.001).




FIG. 2. Multivariate OR of type 2 diabetes according to quintiles of CRP levels stratified by regular aspirin use ( $\geq$ 1/week). *P* for interaction = 0.0002. Adjusted for the same variables as in Table 3.

## DISCUSSION

In this large prospective, nested, case-control study, elevated baseline levels of TNF- $\alpha$ R2, IL-6, and CRP were significantly associated with the risk of developing type 2 diabetes over 10 years of follow-up. These associations were independent of BMI, physical activity, and other conventional diabetes risk factors and were not explained by hyperinsulinemia. Among the three inflammatory markers, CRP had the strongest and most robust relationship with diabetes.

The concept of atherosclerosis as an inflammatory disease is now well established (30). Recent cross-sectional data have demonstrated that elevated levels of CRP

are associated with obesity, insulin resistance, and glucose intolerance (31–33), suggesting that inflammation is also involved in the etiology of type 2 diabetes. However, prospective (or nested case-control) studies evaluating the association between CRP levels and risk of type 2 diabetes have been inconsistent. Several studies have reported a significant positive association, even after adjustment for BMI (9–12). However, in the Insulin Resistance Atherosclerosis Study (14) and the MONICA Augsburg Cohort Study (15), the association of CRP with diabetes disappeared after adjustment for BMI. In the Mexico City Diabetes Study, CRP was an independent predictor of metabolic syndrome and type 2 diabetes in women but not in men (13,34). In a nested case-control study of 71 pairs of diabetes case and control subjects (matched on BMI) in the Pima Indians (16), CRP was not related to risk of diabetes. Our study, with a much larger sample size, found a strong and independent association between CRP levels and type 2 diabetes. The magnitude of association between CRP and diabetes appears to be comparable or even stronger than the association of CRP with coronary heart disease observed in previous studies (30). These data support the hypothesis that low-grade systemic inflammation is a common antecedent for both type 2 diabetes and CVD (35). Consistent with previous results with coronary disease (36), we found that the association between CRP and diabetes was significantly stronger among nonaspirin users than aspirin users, suggesting that the inflammatory role of CRP may be mitigated by aspirin use.

The biological mechanisms through which CRP increases risk of type 2 diabetes are not well understood. CRP is a marker of low-grade inflammation and may have indirect influence on insulin resistance and insulin secretion through altered innate immune response due to heightened systemic inflammation (37,38). Elevated CRP also stimulates endothelial production of E-selectin, ICAM-1, and VCAM-1, important mediators of impaired vascular reactivity, reduced insulin delivery, and increased peripheral insulin resistance (8). Thus, the positive association between CRP and type 2 diabetes may simply reflect underlying endothelial dysfunction and subclinical atherosclerosis. However, our results show that the effects of CRP are independent of levels of E-selectin, and that the combination of higher CRP and E-selectin levels confers the greatest risk of diabetes. Our results did not change after excluding subjects with incident CVD during the follow-up, suggesting that the positive association of diabetes with CRP was unlikely due to subclinical CVD.

The production of CRP is regulated by inflammatory cytokines such as TNF- $\alpha$  and IL-6. Thus, the observed association between CRP levels and diabetes may partly reflect the detrimental effects of cytokines, such as IL-6 and TNF- $\alpha$ , on insulin resistance. Adipose tissue is a major source of endogenous TNF- $\alpha$  production; elevation in levels of TNF- $\alpha$  may be a critical mechanism by which fat cells induce peripheral insulin resistance (39). Elevations in TNF- $\alpha$  levels then reflect both upregulated adipocyte signaling and systemic inflammation (6,40). TNF- $\alpha$  may mediate insulin resistance through indirect effects, including increasing free fatty acid oxidation, stimulation of insulin counterregulatory hormones or cytokines (e.g., IL-6 and CRP), impairment of endothelial function, or

direct inhibitory effects on glucose transporter protein GLUT4, insulin receptor substrates, or glucose-stimulated insulin release by pancreatic  $\beta$ -cells. The results from our study suggest that the effects of TNF- $\alpha$  were accounted for by elevated CRP levels and may be mediated through obesity and elevated production of CRP. Interestingly, the positive association between TNF- $\alpha$ R2 was more evident in nonobese than obese women. This result is consistent with previous findings that TNF- $\alpha$ R2 was more strongly correlated with leptin, insulin, and C-peptide among normal weight than overweight subjects (41).

Consistent with the Women's Health Study (9), we found that the predictive role of IL-6 was weaker than that of CRP, and the association was also accounted for by elevated levels of CRP. Weaker associations of diabetes with TNF- $\alpha$ R2 and IL-6 may result from larger variability and thus greater random misclassification in the measurement. Clinically, CRP is a more stable and integrated measure of low-grade inflammation. Biologically, however, IL-6 may serve an inflammatory role by modulating the production of TNF- $\alpha$ , downregulating the immune response, and upregulating the production of CRP (8).

The strengths of our study include the large sample size, a long follow-up, and detailed measures of diet and lifestyle. Several limitations should be considered, however. First, because all our participants were health professionals, our findings may not directly apply to the general population. Second, diabetes diagnoses were reported by the nurses but confirmed by a validated supplementary questionnaire regarding symptoms, diagnostic tests, and treatment. Our previous study found this confirmation highly accurate compared with medical record review (19). Nonetheless, the findings of our study may only apply to the clinically diagnosed case subjects because it was not feasible to screen the entire cohort. Because our control subjects were not uniformly screened for glucose intolerance, some cases of diabetes may have been undiagnosed. However, when the analyses were restricted to subjects with HbA<sub>1c</sub> <6.0% at baseline or adjusted for baseline HbA<sub>1c</sub> levels, the findings were not appreciably altered, suggesting that bias due to undiagnosed diabetes is unlikely. Third, while direct TNF- $\alpha$  measurement in plasma is not possible using frozen specimens, TNF- $\alpha$  signals through at least two known cell surface receptors, TNF- $\alpha$ R1 (receptor 1) and TNF- $\alpha$ R2 (receptor 2). These receptors are shed into the plasma, and TNF- $\alpha$ R2, a specific marker of TNF- $\alpha$ -related insulin resistance, is easily and reliably measured in frozen plasma (42).

The clinical implications of these data are twofold. First, elevated CRP levels in apparently healthy subjects may help to identify high-risk populations for both type 2 diabetes and CVD. So far, CRP appears to be the most consistent and strongest nonconventional predictor of both conditions. Second, elevated CRP can serve as a common target for lifestyle and therapeutic interventions for these two conditions. Lifestyle interventions such as exercise and weight loss are effective in lowering CRP and other inflammatory markers (43,44) and in preventing type 2 diabetes in subjects with impaired glucose tolerance (45,46). Therapy with statins and ACE inhibitors have been shown to reduce CRP and improve insulin sensitivity and

endothelial function (47,48) and, in secondary analyses of CVD prevention studies, have been associated with a reduced risk of incident type 2 diabetes (49–51). However, whether a reduction in CRP per se will lead to a reduction in risk of type 2 diabetes and CVD remains to be seen in randomized trials.

In conclusion, we found that elevated plasma levels of inflammatory markers, especially CRP, were independent predictors of type 2 diabetes in apparently healthy women. Our findings support the hypothesis that low-grade systemic inflammation is an underlying factor in the pathogenesis of type 2 diabetes. These findings may have important implications for the prevention and treatment of type 2 diabetes. Modification of unhealthy diet and lifestyle factors, fundamental causes of heightened inflammatory response, and type 2 diabetes should remain the cornerstone for the prevention and management of diabetes. Pharmacological agents with anti-inflammatory properties may also have a role in diabetes prevention and treatment.

#### ACKNOWLEDGMENTS

This study was supported by research Grants CA87969 and DK58845 from the National Institutes of Health.

We are indebted to the participants in the Nurses' Health Study for their cooperation and to Jeanne Sparrow for laboratory assistance.

#### REFERENCES

- Wilding JPH: Obesity and nutritional factors in the pathogenesis of type 2 diabetes mellitus. In *Textbook of Diabetes*. 3rd ed. Pickup JC, Ed. Oxford, U.K., Blackwell Science Ltd., 2003, p. 21.21–21.16
- Das UN: Is obesity an inflammatory condition? *Nutrition* 17:953–966, 2001
- Yudkin JS, Stehouwer CD, Emeis JJ, Coppack SW: C-reactive protein in healthy subjects: associations with obesity, insulin resistance, and endothelial dysfunction: a potential role for cytokines originating from adipose tissue? *Arterioscler Thromb Vasc Biol* 19:972–978, 1999
- Hu FB, Stampfer MJ: Is type 2 diabetes mellitus a vascular condition? *Arterioscler Thromb Vasc Biol* 23:1715–1716, 2003
- Hotamisligil GS, Spiegelman BM: Tumor necrosis factor- $\alpha$ : a key component of the obesity-diabetes link. *Diabetes* 43:1271–1278, 1994
- Hotamisligil GS, Arner P, Caro JF, Atkinson RL, Spiegelman BM: Increased adipose tissue expression of tumor necrosis factor- $\alpha$  in human obesity and insulin resistance. *J Clin Invest* 95:2409–2415, 1995
- Caballero AE, Arora S, Saouaf R, Lim SC, Smakowski P, Park JY, King GL, LoGerfo FW, Horton ES, Veves A: Microvascular and macrovascular reactivity is reduced in subjects at risk for type 2 diabetes. *Diabetes* 48:1856–1862, 1999
- Gabay C, Kushner I: Acute-phase proteins and other systemic responses to inflammation *N Engl J Med* 340:448–454, 1999. [published erratum appears in *N Engl J Med* 29:1376, 1999]
- Pradhan AD, Manson JE, Rifai N, Buring JE, Ridker PM: C-reactive protein, interleukin 6, and risk of developing type 2 diabetes mellitus. *JAMA* 286:327–334, 2001
- Barzilay JI, Abraham L, Heckbert SR, Cushman M, Kuller LH, Resnick HE, Tracy RP: The relation of markers of inflammation to the development of glucose disorders in the elderly: the Cardiovascular Health Study. *Diabetes* 50:2384–2389, 2001
- Freeman DJ, Norrie J, Caslake MJ, Gaw A, Ford I, Lowe GD, O'Reilly DS, Packard CJ, Sattar N: C-reactive protein is an independent predictor of risk for the development of diabetes in the West of Scotland Coronary Prevention Study. *Diabetes* 51:1596–1600, 2002
- Spranger J, Kroke A, Mhlig M, Hoffmann K, Bergmann MM, Ristow M, Boeing H, Pfeiffer AF: Inflammatory cytokines and the risk to develop type 2 diabetes: results of the prospective population-based European Prospective Investigation into Cancer and Nutrition (EPIC)-Potsdam Study. *Diabetes* 52:812–817, 2003
- Han TS, Sattar N, Williams K, Gonzalez-Villalpando C, Lean ME, Haffner SM: Prospective study of C-reactive protein in relation to the development of diabetes and metabolic syndrome in the Mexico City Diabetes Study. *Diabetes Care* 25:2016–2021, 2002
- Festa A, D'Agostino R Jr, Tracy RP, Haffner SM: Elevated levels of acute-phase proteins and plasminogen activator inhibitor-1 predict the development of type 2 diabetes: the Insulin Resistance Atherosclerosis Study. *Diabetes* 51:1131–1137, 2002
- Thorand B, Lowel H, Schneider A, Kolb H, Meisinger C, Frohlich M, Koenig W: C-reactive protein as a predictor for incident diabetes mellitus among middle-aged men: results from the MONICA Augsburg cohort study, 1984–1998. *Arch Intern Med* 163:93–99, 2003
- Krakoff J, Funahashi T, Stehouwer CD, Schalkwijk CG, Tanaka S, Matsuzawa Y, Kobes S, Tataranni PA, Hanson RL, Knowler WC, Lindsay RS: Inflammatory markers, Adiponectin, and risk of type 2 diabetes in the Pima Indian. *Diabetes Care* 26:1745–1751, 2003
- National Diabetes Data Group: Classification and diagnosis of diabetes mellitus and other categories of glucose intolerance. *Diabetes* 28:1039–1057, 1979
- Expert Committee on the Diagnosis and Classification of Diabetes Mellitus: Report of the Expert Committee on the Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 20:1183–1194, 1997
- Manson JE, Colditz GA, Stampfer MJ, Willett WC, Krolewski AS, Rosner B, Arky RA, Speizer FE, Hennekens CH: A prospective study of maturity-onset diabetes mellitus and risk of coronary heart disease and stroke in women. *Arch Intern Med* 151:1141–1147, 1991
- Field AE, Coakley EH, Must A, Spadano JL, Laird N, Dietz WH, Rimm E, Colditz GA: Impact of overweight on the risk of developing common chronic diseases during a 10-year period. *Arch Intern Med* 161:1581–1586, 2001
- Manson JE, Willett WC, Stampfer MJ, Colditz GA, Hunter DJ, Hankinson SE, Hennekens CH, Speizer FE: Body weight and mortality among women. *N Engl J Med* 333:677–685, 1995
- Carey VJ, Walters EE, Colditz GA, Solomon CG, Willett WC, Rosner BA, Speizer FE, Manson JE: Body fat distribution and risk of non-insulin-dependent diabetes mellitus in women: the Nurses' Health Study. *Am J Epidemiol* 145:614–619, 1997
- Hu FB, Sigal RJ, Rich-Edwards JW, Colditz GA, Solomon CG, Willett WC, Speizer FE, Manson JE: Walking compared with vigorous physical activity and risk of type 2 diabetes in women: a prospective study. *JAMA* 282:1433–1439, 1999
- Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, Hennekens CH, Speizer FE: Reproducibility and validity of a semi-quantitative food frequency questionnaire. *Am J Epidemiol* 122:51–65, 1985
- Hu FB, Manson JE, Stampfer MJ, Colditz G, Liu S, Solomon CG, Willett WC: Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *N Engl J Med* 345:790–797, 2001
- Hankinson SE, London SJ, Chute CG, Barbieri RL, Jones L, Kaplan LA, Sacks FM, Stampfer MJ: Effect of transport conditions on the stability of biochemical markers in blood. *Clin Chem* 35:2313–2316, 1989
- SAS Institute Inc: *SAS/STAT User's Guide*. V. 6, 4th ed., vol. 1. Cary, NC, SAS Institute Inc., 1989
- Duncan BB, Schmidt MI, Pankow JS, Ballantyne CM, Couper D, Vigo A, Hoogeveen R, Folsom AR, Heiss G: Low-grade systemic inflammation and the development of type 2 diabetes: the Atherosclerosis Risk in Communities Study. *Diabetes* 52:1799–1805, 2003
- Meigs JB, Hu FB, Manson JE: Endothelial dysfunction predicts development of type 2 diabetes (Abstract). *Diabetes* 52 (Suppl. 1):A58, 2003
- Ridker PM: Clinical application of C-reactive protein for cardiovascular disease detection and prevention. *Circulation* 107:363–369, 2003
- Visser M, Bouter LM, McQuillan GM, Wener MH, Harris TB: Elevated C-reactive protein levels in overweight and obese adults. *JAMA* 282:2131–2135, 1999
- Festa A, D'Agostino R Jr, Howard G, Mykkanen L, Tracy RP, Haffner SM: Chronic subclinical inflammation as part of the insulin resistance syndrome: the Insulin Resistance Atherosclerosis Study (IRAS). *Circulation* 102:42–47, 2000
- Pradhan AD, Cook NR, Buring JE, Manson JE, Ridker PM: C-reactive protein is independently associated with fasting insulin in nondiabetic women. *Arterioscler Thromb Vasc Biol* 23:650–655, 2003
- Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33): UK Prospective Diabetes Study (UKPDS) Group. *Lancet* 352:837–853, 1998
- Pradhan AD, Ridker PM: Do atherosclerosis and type 2 diabetes share a common inflammatory basis? *Eur Heart J* 23:831–834, 2002
- Ridker PM, Cushman M, Stampfer MJ, Tracy RP, Hennekens CH: Inflam-

- mation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med* 336:973-979, 1997
37. Pickup JC, Mattock MB, Chusney GD, Burt D: NIDDM as a disease of the innate immune system: association of acute-phase reactants and interleukin-6 with metabolic syndrome X. *Diabetologia* 40:1286-1292, 1997
  38. Pickup JC, Crook MA: Is type II diabetes mellitus a disease of the innate immune system? *Diabetologia* 41:1241-1248, 1998
  39. Hotamisligil GS, Shargill NS, Spiegelman BM: Adipose expression of tumor necrosis factor- $\alpha$ : direct role in obesity-linked insulin resistance. *Science* 259:87-91, 1993
  40. Saghizadeh M, Ong JM, Garvey WT, Henry RR, Kern PA: The expression of TNF  $\alpha$  by human muscle: relationship to insulin resistance. *J Clin Invest* 97:1111-1116, 1996
  41. Chu NF, Spiegelman D, Rifai N, Hotamisligil GS, Rimm EB: Glycemic status and soluble tumor necrosis factor receptor levels in relation to plasma leptin concentrations among normal weight and overweight US men. *Int J Obes Relat Metab Disord* 24:1085-1092, 2000
  42. Chu NF, Spiegelman D, Hotamisligil GS, Rifai N, Stampfer M, Rimm EB: Plasma insulin, leptin, and soluble TNF receptors levels in relation to obesity-related atherogenic and thrombogenic cardiovascular disease risk factors among men. *Atherosclerosis* 157:495-503, 2001
  43. Esposito K, Pontillo A, Di Palo C, Giugliano G, Masella M, Marfella R, Giugliano D: Effect of weight loss and lifestyle changes on vascular inflammatory markers in obese women: a randomized trial. *JAMA* 289:1799-1804, 2003
  44. Manns PJ, Williams DP, Snow CM, Wander RC: Physical activity, body fat, and serum C-reactive protein in postmenopausal women with and without hormone replacement. *Am J Human Biol* 15:91-100, 2003
  45. Tuomilehto J, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, Ilanne-Parikka P, Keinanen-Kiukkaanniemi S, Laakso M, Louheranta A, Rastas M, Salminen V, Uusitupa M: Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med* 344:1343-1350, 2001
  46. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM: Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med* 346:393-403, 2002
  47. Tan KC, Chow WS, Tam SC, Ai VH, Lam CH, Lam KS: Atorvastatin lowers C-reactive protein and improves endothelium-dependent vasodilation in type 2 diabetes mellitus. *J Clin Endocrinol Metab* 87:563-568, 2002
  48. O'Driscoll G, Green D, Maiorana A, Stanton K, Colreavy F, Taylor R: Improvement in endothelial function by angiotensin-converting enzyme inhibition in non-insulin-dependent diabetes mellitus. *J Am Coll Cardiol* 33:1506-1511, 1999
  49. Freeman DJ, Norrie J, Sattar N, Neely RD, Cobbe SM, Ford I, Isles C, Lorimer AR, Macfarlane PW, McKillop JH, Packard CJ, Shepherd J, Gaw A: Pravastatin and the development of diabetes mellitus: evidence for a protective treatment effect in the West of Scotland Coronary Prevention Study. *Circulation* 103:357-362, 2001
  50. Yusuf S, Gerstein H, Hoogwerf B, Pogue J, Bosch J, Wolfenbittel BH, Zinman B: Ramipril and the development of diabetes. *JAMA* 286:1882-1885, 2001
  51. Vermees E, Ducharme A, Bourassa MG, Lessard M, White M, Tardif JC: Enalapril reduces the incidence of diabetes in patients with chronic heart failure: insight from the Studies Of Left Ventricular Dysfunction (SOLVD). *Circulation* 107:1291-1296, 2003